

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 1

Question:

A particle P is moving in a straight line. Initially P is moving through a point O with speed 4 m s^{-1} . At time t seconds after passing through O the acceleration of P is $3e^{-0.25t} \text{ m s}^{-2}$ in the direction OP . Find the velocity of the particle at time t seconds.

Solution:

$$\begin{aligned}v &= \int a \, dt = \int 3e^{-0.25t} \, dt \\ &= -12e^{-0.25t} + A\end{aligned}$$

When $t = 0, v = 4$

$$4 = -12 + A \Rightarrow A = 16$$

$$v = 16 - 12e^{-0.25t}$$

The velocity of the particle at time t seconds is $(16 - 12e^{-0.25t}) \text{ m s}^{-1}$.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 2

Question:

A particle P is moving along the x -axis in the direction of x increasing. At time t seconds, the velocity of P is $(t \sin t) \text{ m s}^{-1}$. When $t = 0$, P is at the origin. Show that when $t = \frac{\pi}{2}$, P is 1 metre from O .

Solution:

$$x = \int v \, dt = \int t \sin t \, dt$$

Using integration by parts

$$x = -t \cos t + \int \cos t \, dt$$

$$= -t \cos t + \sin t + A$$

$$t = 0, x = 0$$

$$\text{When } 0 = 0 + 0 + A \Rightarrow A = 0$$

$$x = -t \cos t + \sin t$$

$$\text{When } t = \frac{\pi}{2}$$

$$x = -\frac{\pi}{2} \cos \frac{\pi}{2} + \sin \frac{\pi}{2} = 1$$

Hence P is one metre from O , as required.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 3

Question:

At time t seconds the velocity, v m s⁻¹, of a particle moving in a straight line is given

$$\text{by } v = \frac{4}{3+2t^0} \quad t \geq 0.$$

When $t = 0$, the particle is at a point A . When $t = 3$, the particle is at the point B .

Find the distance between A and B .

Solution:

$$s = \int v \, dt = \int \frac{4}{3+2t} \, dt = 2\ln(3+2t) + C \quad \text{where } s \text{ is the displacement from point } A.$$

$$\text{When } t = 0, s = 0$$

$$0 = 2\ln 3 + C \Rightarrow C = -2\ln 3$$

$$s = 2\ln(3+2t) - 2\ln 3 = 2\ln\left(\frac{3+2t}{3}\right)$$

$$\text{When } t = 3$$

$$s = 2\ln\left(\frac{3+6}{3}\right) = 2\ln 3$$

$$AB = 2\ln 3 \text{ m}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 4

Question:

A particle P is moving along the x -axis in the positive direction. At time t seconds the acceleration of P is $4e^{\frac{1}{2}t}$ m s⁻² in the positive direction. When $t = 0$, P is at rest. Find the distance P moves in the interval $0 \leq t \leq 2$. Give your answer to 3 significant figures.

Solution:

$$v = \int a \, dt = \int 4e^{\frac{1}{2}t} \, dt = 8e^{\frac{1}{2}t} + A$$

When $t = 0$, $v = 0$

$$0 = 8 + A \Rightarrow A = -8$$

$$v = 8e^{\frac{1}{2}t} - 8$$

The distance moved in the interval $0 \leq t \leq 2$ is given by

$$s = \int v \, dt = \int_0^2 \left(8e^{\frac{1}{2}t} - 8 \right) dt$$

$$= \left[16e^{\frac{1}{2}t} - 8t \right]_0^2 = (16e^1 - 16) - 16$$

$$= 16e - 32 \approx 11.5$$

The distance moved is 11.5 m (3 s.f.).

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 5

Question:

A particle P is moving along the x -axis. At time t seconds the displacement of P from O is x m and the velocity of P is $(4 \cos 3t \text{ m s}^{-1})$, both measured in the direction Ox .

When $t = 0$ the particle P is at the origin O . Find

- the magnitude of the acceleration when $t = \frac{\pi}{12}$,
- x in terms of t ,
- the smallest positive value of t for which P is at O .

Solution:

$$\text{a } a = \frac{dv}{dt} = -12 \sin 3t$$

$$\text{When } t = \frac{\pi}{12}$$

$$a = -12 \sin \frac{\pi}{4} = -12 \times \frac{1}{\sqrt{2}} = -6\sqrt{2}$$

The magnitude of the acceleration when $t = \frac{\pi}{12}$ is $6\sqrt{2} \text{ m s}^{-2}$.

$$\text{b } x = \int v \, dt = \int 4 \cos 3t \, dt = \frac{4}{3} \sin 3t + A$$

$$t = 0, x = 0$$

$$\text{When } 0 = \frac{4}{3} \times 0 + A \Rightarrow A = 0$$

$$x = \frac{4}{3} \sin 3t$$

c When P is at O , $x = 0$

$$x = \frac{4}{3} \sin 3t = 0 \Rightarrow \sin 3t = 0$$

The smallest positive value of t is given by

$$3t = \pi \Rightarrow t = \frac{\pi}{3}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 6

Question:

A particle P is moving along a straight line. Initially P is at rest. At time t seconds P has velocity v m s⁻¹ and acceleration α m s⁻² where

$$\alpha = \frac{6t}{(2+t^2)^2} \quad t \geq 0.$$

Find v in terms of t .

Solution:

$$v = \int \alpha \, dt = \int \frac{6t}{(2+t^2)^2} \, dt$$

Let $u = 2 + t^2$, then $\frac{du}{dt} = 2t$

$$\begin{aligned} v &= \int \frac{6t}{(2+t^2)^2} \, dt = \int \frac{3}{(2+t^2)^2} \times 2t \, dt \\ &= \int \frac{3}{u^2} \, du = \int 3u^{-2} \, du \\ &= \frac{3u^{-1}}{-1} + A = A - \frac{3}{u} \\ &= A - \frac{3}{2+t^2} \end{aligned}$$

When $t = 0, v = 0$

$$0 = A - \frac{3}{2} \Rightarrow A = \frac{3}{2}$$

$$v = \frac{3}{2} - \frac{3}{2+t^2}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 7

Question:


A particle P is moving along the x -axis. At time t seconds the velocity of P is $v \text{ m s}^{-1}$ in the direction of x increasing, where

$$v = \begin{cases} 4, & 0 \leq t \leq 3 \\ 5 - \frac{3}{t}, & 3 < t \leq 6 \end{cases}$$

When $t = 0$, P is at the origin O .

- Sketch a velocity-time graph to illustrate the motion of P in the interval $0 \leq t \leq 6$.
- Find the distance of P from O when $t = 6$.

Solution:



- The distance moved in the first three seconds is represented by the area labelled ①.

Let this area be A_1 . $A_1 = 3 \times 4 = 12$

The distance travelled in the next three seconds is represented by the area labelled ②.

Let this area be A_2 .

$$\begin{aligned} A_2 &= \int_3^6 \left(5 - \frac{3}{t} \right) dt \\ &= [5t - 3 \ln t]_3^6 = (30 - 3 \ln 6) - (15 - 3 \ln 3) \\ &= 15 - 3 \ln 2 \end{aligned}$$

The distance of P from O when $t = 6$ is $(12 + 15 - 3 \ln 2) \text{ m} = (27 - 3 \ln 2) \text{ m}$.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 8

Question:

A particle P is moving in a straight line with acceleration $\left(\sin \frac{1}{2}t\right) \text{ m s}^{-2}$ at time

t seconds, $t \geq 0$. The particle is initially at rest at a point O . Find

- the speed of P when $t = 2\pi$,
- the distance of P from O when $t = \frac{\pi}{2}$.

Solution:

$$\text{a } v = \int a \, dt = \int \sin \frac{1}{2}t \, dt = -2 \cos \frac{1}{2}t + A$$

When $t = 0, v = 0$

$$0 = -2 + A \Rightarrow A = 2$$

$$v = 2 - 2 \cos \frac{1}{2}t$$

When $t = 2\pi$

$$v = 2 - 2 \cos \pi = 2 - (2 \times -1) = 4$$

The speed of P when $t = 2\pi$ is 4 m s^{-1} .

$$\text{b } x = \int v \, dt = \int \left(2 - 2 \cos \frac{1}{2}t\right) dt = 2t - 4 \sin \frac{1}{2}t + B$$

When $t = 0, x = 0$

$$0 = 0 - 0 + B \Rightarrow B = 0$$

$$x = 2t - 4 \sin \frac{1}{2}t$$

When $t = \frac{\pi}{2}$

$$x = 2 \times \frac{\pi}{2} - 4 \sin \frac{\pi}{4} = \pi - 4 \times \frac{1}{\sqrt{2}} = \pi - 2\sqrt{2}$$

The distance of P from O when $t = \frac{\pi}{2}$ is $(\pi - 2\sqrt{2}) \text{ m}$.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 9

Question:

A particle P is moving along the x -axis. At time t seconds P has velocity v m s⁻¹ in the direction x increasing and an acceleration of magnitude $4e^{0.2t}$ m s⁻² in the direction x decreasing. When $t = 0$, P is moving through the origin with velocity 20 m s⁻¹ in the direction x increasing. Find

- v in terms of t ,
- the maximum value of x attained by P during its motion.

Solution:

$$\text{a } v = \int a \, dt = \int -4e^{0.2t} \, dt = -20e^{0.2t} + A$$

$$\text{When } t = 0, v = 20$$

$$20 = -20 + A \Rightarrow A = 40$$

$$v = 40 - 20e^{0.2t}$$

$$\text{b } x = \int v \, dt = \int (40 - 20e^{0.2t}) \, dt = 40t - 100e^{0.2t} + B$$

$$\text{When } t = 0, x = 0$$

$$0 = 0 - 100 + B \Rightarrow B = 100$$

$$x = 40t - 100e^{0.2t} + 100$$

The maximum value of x occurs when

$$\frac{dx}{dt} = v = 40 - 20e^{0.2t} = 0$$

$$e^{0.2t} = 2$$

$$0.2t = \ln 2$$

$$t = 5 \ln 2$$

The maximum value of x is given by

$$x = 40 \times 5 \ln 2 - 100 \times 2 + 100 = 200 \ln 2 - 100$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 10

Question:

A car is travelling along a straight road. As it passes a sign S , the driver applies the brakes. The car is modelled as a particle. At time t seconds the car is x m from S and

its velocity, v m s⁻¹, is modelled by the equation $v = \frac{3200}{c + dt}$ where c and d are

constants.

Given that when $t = 0$, the speed of the car is 40 m s⁻¹ and its deceleration is

0.5 m s⁻², find

- the value of c and the value of d ,
- x in terms of t .

Solution:

$$\text{a } v = \frac{3200}{c + dt}$$

When $t = 0$, $v = 40$

$$40 = \frac{3200}{c} \Rightarrow c = 80$$

$$v = \frac{3200}{80 + dt} = 3200(80 + dt)^{-1}$$

$$a = \frac{dv}{dt} = -3200d(80 + dt)^{-2} = -\frac{3200d}{(80 + dt)^2}$$

When $t = 0$, $a = -0.5$

$$-\frac{3200d}{80^2} = -0.5 \Rightarrow d = \frac{0.5 \times 80^2}{3200} = 1$$

$$c = 80, d = 1$$

$$\text{b } x = \int v \, dt = \int \frac{3200}{80 + t} \, dt = 3200 \ln(80 + t) + A$$

When $t = 0$, $x = 0$

$$0 = 3200 \ln 80 + A \Rightarrow A = -3200 \ln 80$$

$$x = 3200 \ln(80 + t) - 3200 \ln 80 = 3200 \ln \left(\frac{80 + t}{80} \right)$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 11

Question:

A particle P is moving along a straight line. When $t=0$, P is passing through a point

A . At time t seconds after passing through A the velocity, v m s⁻¹, of P is given by

$$v = e^{2t} - 11e^t + 15t.$$

Find

- the values of t for which the acceleration is zero,
- the distance of P from A when $t = \ln 3$.

Solution:

$$\text{a } a = \frac{dv}{dt} = 2e^{2t} - 11e^t + 15 = 0$$

$$(2e^t - 5)(2e^t - 3) = 0$$

$$e^t = 2.5, 3$$

$$t = \ln 2.5, \ln 3$$

$$\text{b } x = \int v \, dt = \int (e^{2t} - 11e^t + 15t) \, dt = \frac{e^{2t}}{2} - 11e^t + \frac{15t^2}{2} + A$$

When $t = 0$, $x = 0$

$$0 = \frac{1}{2} - 11 + 0 + A \Rightarrow A = \frac{21}{2}$$

$$x = \frac{e^{2t}}{2} - 11e^t + \frac{15t^2}{2} + \frac{21}{2}$$

When $t = \ln 3$

$$\begin{aligned} x &= \frac{e^{2\ln 3}}{2} - 11e^{\ln 3} + \frac{15(\ln 3)^2}{2} + \frac{21}{2} \\ &= \frac{9}{2} - 33 + \frac{15(\ln 3)^2}{2} + \frac{21}{2} = \frac{15(\ln 3)^2}{2} - 18 \approx -8.95 \end{aligned}$$

As distance is a positive quantity, the required distance is $\left(18 - \frac{15(\ln 3)^2}{2}\right)$ m.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise A, Question 12

Question:

A particle P moves along a straight line. At time t seconds (where $t \geq 0$) the velocity of P is $[2t + \ln(t+2)] \text{ m s}^{-1}$. Find

- the value of t for which the acceleration has magnitude 2.2 m s^{-2} .
- the distance moved by P in the interval $1 \leq t \leq 4$.

Solution:

$$\text{a } a = \frac{dv}{dt} = 2 + \frac{1}{t+2} = 2.2$$

$$\frac{1}{t+2} = 0.2 \Rightarrow t+2 = 5 \Rightarrow t = 3$$

$$\text{b } x = \int v \, dt = \int_1^4 (2t + \ln(t+2)) \, dt$$

Using integration by parts

$$\int \ln(t+2) \, dt = \int 1 \cdot \ln(t+2) \, dt$$

$$= t \ln(t+2) - \int \frac{t}{t+2} \, dt = t \ln(t+2) - \int \left(1 - \frac{2}{t+2}\right) dt$$

$$= t \ln(t+2) - t + 2 \ln(t+2) = (t+2) \ln(t+2) - t$$

$$\begin{aligned} \text{Hence } x &= \left[(t+2) \ln(t+2) - t \right]_1^4 \\ &= (16 + 6 \ln 6 - 4) - (1 + 3 \ln 3 - 1) \\ &= 12 + 6 \ln 6 - 3 \ln 3 = 12 + 3 \ln 6^2 - 3 \ln 3 \\ &= 12 + 3 \ln \left(\frac{36}{3} \right) = 12 + 3 \ln 12 \end{aligned}$$

The distance moved by P in the interval $1 \leq t \leq 4$ is $(12 + 3 \ln 12) \text{ m}$.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 1

Question:

A particle P moves along the x -axis. At time $t=0$, P passes through the origin O with velocity 5 m s^{-1} in the direction of x increasing. At time t seconds, the velocity of P is $v \text{ m s}^{-1}$ and $OP = x \text{ m}$. The acceleration of P is $\left(2 + \frac{1}{2}x\right) \text{ m s}^{-2}$, measured in the positive x direction. Find v^2 in terms of x .

Solution:

$$a = 2 + \frac{1}{2}x$$

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = 2 + \frac{1}{2}x$$

$$\frac{1}{2}v^2 = \int\left(2 + \frac{1}{2}x\right) dx$$

$$= 2x + \frac{x^2}{4} + A$$

At $x = 0, v = 5$

$$\frac{1}{2} \times 25 = 0 + 0 + A \Rightarrow A = \frac{25}{2}$$

$$\frac{1}{2}v^2 = 2x + \frac{x^2}{4} + \frac{25}{2}$$

$$v^2 = \frac{x^2}{2} + 4x + 25$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 2

Question:

A particle P moves along a straight line. When its displacement from a fixed point O on the line is x m and its velocity is v m s⁻¹, the deceleration of P is $4x$ m s⁻². At $x = 2$, $v = 8$. Find v in terms of x .

Solution:

$$a = -4x$$

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -4x$$

$$\frac{1}{2}v^2 = \int (-4x) dx$$
$$= -2x^2 + A$$

$$\text{At } x = 2, v = 8$$

$$\frac{1}{2} \times 64 = -8 + A \Rightarrow A = 40$$

$$\frac{1}{2}v^2 = -2x^2 + 40$$

$$v^2 = 80 - 4x^2$$

$$v = \pm\sqrt{(80 - 4x^2)}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 3

Question:

A particle P is moving along the x -axis in the direction of x increasing. At $OP = x \text{ m} (x > 0)$, the velocity of P is $v \text{ m s}^{-1}$ and its acceleration is of magnitude $\frac{4}{x^2} \text{ m s}^{-2}$ in the direction of x increasing. Given that at $x = 2, v = 6$ find the value of x for which P is instantaneously at rest.

Solution:

$$a = \frac{4}{x^2}$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = \frac{4}{x^2}$$

$$\frac{1}{2} v^2 = \int (4x^{-2}) dx$$

$$= \frac{4x^{-1}}{-1} + A = A - \frac{4}{x}$$

At $x = 2, v = 6$

$$\frac{1}{2} \times 36 = A - 2 \Rightarrow A = 20$$

$$\frac{1}{2} v^2 = 20 - \frac{4}{x}$$

When $v = 0$

$$0 = 20 - \frac{4}{x} \Rightarrow x = \frac{4}{20} = \frac{1}{5}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 4

Question:

A particle P moves along a straight line. When its displacement from a fixed point O on the line is x m and its velocity is v m s⁻¹, the acceleration of P is of magnitude $25x$ m s⁻² and is directed towards O . At $x = 0, v = 40$. In its motion P is instantaneously at rest at two points A and B . Find the distance between A and B .

Solution:

$$a = -25x$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = -25x$$

$$\frac{1}{2} v^2 = \int (-25x) dx$$

$$= -\frac{25}{2} x^2 + A$$

At $x = 0, v = 40$

$$\frac{1}{2} \times 1600 = -0 + A \Rightarrow A = 800$$

$$\frac{1}{2} v^2 = -\frac{25}{2} x^2 + 800$$

When $v = 0$

$$0 = -\frac{25}{2} x^2 + 800 \Rightarrow x^2 = 64 \Rightarrow x = \pm 8$$

$$AB = 16 \text{ m}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 5

Question:

A particle P is moving along the x -axis. At $OP = x$ m, the velocity of P is v m s⁻¹ and its acceleration is of magnitude kx^2 m s⁻², where k is a positive constant, in the direction of x decreasing. At $x = 0$, $v = 16$. The particle is instantaneously at rest at $x = 20$. Find

- the value of k ,
- the velocity of P when $x = 10$.

Solution:

$$\begin{aligned} \text{a } a &= -kx^2 \\ \frac{d}{dx} \left(\frac{1}{2}v^2 \right) &= -kx^2 \\ \frac{1}{2}v^2 &= \int (-kx^2) dx \\ &= -\frac{kx^3}{3} + A \end{aligned}$$

At $x = 0$, $v = 16$

$$\frac{1}{2} \times 256 = -0 + A \Rightarrow A = 128$$

$$\frac{1}{2}v^2 = -\frac{kx^3}{3} + 128$$

When $v = 0$, $x = 20$

$$0 = -\frac{8000k}{3} + 128$$

$$k = \frac{3 \times 128}{8000} = \frac{6}{125}$$

$$\text{b } \frac{1}{2}v^2 = -\frac{6}{125} \times \frac{x^3}{3} + 128$$

$$\begin{aligned} v^2 &= 256 - \frac{4}{125}x^3 \\ x &= 10 \end{aligned}$$

$$\text{When } v^2 = 256 - \frac{4}{125} \times 1000 = 224$$

$$v = \pm \sqrt{224} = \pm 4\sqrt{14}$$

The velocity of P when $x = 10$ is $\pm 4\sqrt{14}$ m s⁻¹ as the particle will pass through this position in both directions.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 6

Question:

A particle P is moving along the x -axis in the direction of x increasing. At $OP = x$ m, the velocity of P is v m s^{-1} and its acceleration is of magnitude $8x^3$ m s^{-2} in the direction PO . At $x = 2$, $v = 32$. Find the value of x for which $v = 8$.

Solution:

$$a = -8x^3$$

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -8x^3$$

$$\frac{1}{2}v^2 = \int (-8x^3) dx$$

$$= -2x^4 + A$$

At $x = 2, v = 32$

$$\frac{1}{2} \times 1024 = A - 32 \Rightarrow A = 544$$

$$\frac{1}{2}v^2 = 544 - 2x^4$$

$$v^2 = 1088 - 4x^4$$

When $v = 8$

$$64 = 1088 - 4x^4 \Rightarrow x^4 = 256$$

$$x = 256^{\frac{1}{4}} = 4$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 7

Question:

A particle P is moving along the x -axis. When the displacement of P from the origin O is x m, the velocity of P is v m s⁻¹ and its acceleration is $6 \sin \frac{x}{3}$ m s⁻². At $x = 0$,

$v = 4$. Find

- v^2 in terms of x ,
- the greatest possible speed of P .

Solution:

$$\text{a } a = 6 \sin \frac{x}{3}$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 6 \sin \frac{x}{3}$$

$$\begin{aligned} \frac{1}{2} v^2 &= \int \left(6 \sin \frac{x}{3} \right) dx \\ &= -18 \cos \frac{x}{3} + A \end{aligned}$$

At $x = 0, v = 4$

$$\frac{1}{2} \times 16 = -18 + A \Rightarrow A = 26$$

$$\frac{1}{2} v^2 = -18 \cos \frac{x}{3} + 26$$

$$v^2 = 52 - 36 \cos \frac{x}{3}$$

- The greatest value of v^2 occurs when $\cos \frac{x}{3} = -1$.

The greatest value of v^2 is given by

$$v^2 = 52 + 36 = 88$$

$$v = \pm \sqrt{88} = \pm 2\sqrt{22}$$

The greatest possible speed of P is $2\sqrt{22}$ m s⁻¹ (≈ 9.38 m s⁻¹).

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 8

Question:

A particle P is moving along the x -axis. At $x = 0$, the velocity of P is 2 m s^{-1} in the direction of x increasing. At $OP = x \text{ m}$, the velocity of P is $v \text{ m s}^{-1}$ and its acceleration is $(2 + 3e^{-x}) \text{ m s}^{-2}$. Find the velocity of P at $x = 3$. Give your answer to 3 significant figures.

Solution:

$$a = 2 + 3e^{-x}$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 2 + 3e^{-x}$$

$$\begin{aligned} \frac{1}{2} v^2 &= \int (2 + 3e^{-x}) dx \\ &= 2x - 3e^{-x} + A \end{aligned}$$

$$\text{At } x = 0, v = 2$$

$$\frac{1}{2} \times 4 = 0 - 3 + A \Rightarrow A = 5$$

$$\begin{aligned} \frac{1}{2} v^2 &= 2x - 3e^{-x} + 5 \\ v^2 &= 4x - 6e^{-x} + 10 \end{aligned}$$

$$\text{At } x = 3$$

$$v^2 = 12 - 6e^{-3} + 10 = 21.701\dots$$

$$v = \sqrt{21.701\dots} = 4.658\dots$$

The velocity of P at $x = 3$ is 4.66 m s^{-1} (3 s.f.), in the direction of x increasing.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 9

Question:

A particle P moves away from the origin O along the positive x -axis. The acceleration of P is of magnitude $\frac{4}{2x+1} \text{ m s}^{-2}$, where $OP = x \text{ m}$, directed towards O . Given that

the speed of P at O is 4 m s^{-1} , find

- the speed of P at $x = 10$,
- the value of x at which P is instantaneously at rest.

Give your answers to 3 significant figures.

Solution:

$$\text{a } a = -\frac{4}{2x+1}$$

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -\frac{4}{2x+1}$$

$$\frac{1}{2}v^2 = \int \left(-\frac{4}{2x+1}\right) dx$$

$$= -2\ln(2x+1) + A$$

$$\text{At } x = 0, v = 4$$

$$\frac{1}{2} \times 16 = -0 + A \Rightarrow A = 8$$

$$\frac{1}{2}v^2 = -2\ln(2x+1) + 8$$

$$v^2 = 16 - 4\ln(2x+1)$$

$$\text{At } x = 10$$

$$v^2 = 16 - 4\ln 21 = 3.821910\dots$$

$$v = 1.954\dots$$

The speed of P at $x = 10$ is 1.95 m s^{-1} (3 s.f.).

- When $v = 0$

$$0 = 16 - 4\ln(2x+1) \Rightarrow \ln(2x+1) = 4$$

$$2x+1 = e^4 \Rightarrow x = \frac{e^4 - 1}{2} = 26.799\dots = 26.8 \text{ (3 s.f.)}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 10

Question:

A particle P is moving along the positive x -axis. At $OP = x$ m, the velocity of P is

v m s⁻¹ and its acceleration is $\left(x - \frac{4}{x^3}\right)$ m s⁻². The particle starts from the position

where $x = 1$ with velocity 3 m s⁻¹ in the direction of x increasing. Find

- v in terms of x ,
- the least speed of P during its motion.

Solution:

$$\text{a } a = x - \frac{4}{x^3}$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = x - \frac{4}{x^3}$$

$$\begin{aligned} \frac{1}{2} v^2 &= \int (x - 4x^{-3}) dx \\ &= \frac{x^2}{2} - \frac{4x^{-2}}{-2} + A = \frac{x^2}{2} + \frac{2}{x^2} + A \end{aligned}$$

At $x = 1, v = 3$

$$\frac{1}{2} \times 9 = \frac{1}{2} + 2 + A \Rightarrow A = 2$$

$$\frac{1}{2} v^2 = \frac{x^2}{2} + \frac{2}{x^2} + 2$$

$$v^2 = x^2 + 4 + \frac{4}{x^2} = \left(x + \frac{2}{x} \right)^2$$

$$v = x + \frac{2}{x}$$

- The minimum value of v occurs when $\frac{dv}{dx} = a = 0$

$$x - \frac{4}{x^3} = 0 \Rightarrow x^4 = 4 \Rightarrow x = \sqrt{2} \quad (\text{as } P \text{ moves on the positive } x\text{-axis, } x > 0)$$

At $x = \sqrt{2}$

$$v = \sqrt{2} + \frac{2}{\sqrt{2}} = 2\sqrt{2}$$

The least speed of P during its motion is $2\sqrt{2}$ m s⁻¹.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 11

Question:

A particle P is moving along the x -axis. Initially P is at the origin O moving with velocity 15 m s^{-1} in the direction of x increasing. When the displacement of P from O is $x \text{ m}$, its acceleration is of magnitude $\left(10 + \frac{1}{4}x\right) \text{ m s}^{-2}$ directed towards O . Find the distance P moves before first coming to instantaneous rest.

Solution:

$$a = -\left(10 + \frac{1}{4}x\right)$$

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -10 - \frac{1}{4}x$$

$$\frac{1}{2}v^2 = \int\left(-10 - \frac{1}{4}x\right)dx$$

$$= -10x - \frac{x^2}{8} + A$$

At $x = 0, v = 15$

$$\frac{1}{2} \times 225 = -0 - 0 + A \Rightarrow A = \frac{225}{2}$$

$$\frac{1}{2}v^2 = -10x - \frac{x^2}{8} + \frac{225}{2}$$

$$v^2 = 225 - 20x - \frac{x^2}{4} = -\frac{x^2 + 80x - 900}{4} = -\frac{(x+90)(x-10)}{4}$$

$$v = 0 \Rightarrow x = 10, -90$$

As P is initially moving in the direction of x increasing, it reaches $x = 10$ before $x = -90$. The distance P moves before first coming to instantaneous rest is 10 m.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise B, Question 12

Question:

A particle P is moving along the x -axis. At time t seconds, P is x m from O , has velocity v m s⁻¹ and acceleration of magnitude $6x^{\frac{1}{3}}$ ms⁻² in the direction of x increasing. When $t = 0$, $x = 8$ and $v = 12$. Find

- v in terms of x ,
- x in terms of t .

Solution:

a $a = 6x^{\frac{1}{3}}$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 6x^{\frac{1}{3}}$$

$$\frac{1}{2} v^2 = \int 6x^{\frac{1}{3}} dx = \frac{6x^{\frac{4}{3}}}{\frac{4}{3}} + A = \frac{9}{2} x^{\frac{4}{3}} + A$$

$$v^2 = 9x^{\frac{4}{3}} + B, \text{ where } B = 2A$$

At $x = 8, v = 12$

$$144 = 9 \times 16 + B \Rightarrow B = 0$$

$$v^2 = 9x^{\frac{4}{3}}$$

$$v = 3x^{\frac{2}{3}}$$

b $v = \frac{dx}{dt} = 3x^{\frac{2}{3}}$

Separating the variables and integrating

$$\int x^{-\frac{2}{3}} dx = \int 3 dt$$

$$3x^{\frac{1}{3}} = 3t + C$$

When $t = 0, x = 8$

$$3 \times 2 = 0 + C \Rightarrow C = 6$$

$$3x^{\frac{1}{3}} = 3t + 6$$

$$x^{\frac{1}{3}} = t + 2$$

$$x = (t + 2)^3$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 1

Question:

A particle P moves along a straight line. When the displacement of P from a fixed point on the line is x m, its velocity is v m s⁻¹ and its acceleration is of magnitude

$\frac{6}{x^2}$ m s⁻² in the direction of x increasing. At $x = 3$, $v = 4$.

Find v in terms of x .

Solution:

$$a = \frac{6}{x^2} = 6x^{-2}$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 6x^{-2}$$

$$\frac{1}{2} v^2 = \int 6x^{-2} dx = \frac{6x^{-1}}{-1} + A$$

$$\frac{1}{2} v^2 = A - \frac{6}{x}$$

At $x = 3$, $v = 4$

$$\frac{1}{2} \times 16 = A - 2 \Rightarrow A = 10$$

$$\frac{1}{2} v^2 = 10 - \frac{6}{x}$$

$$v^2 = 20 - \frac{12}{x}$$

$$v = \sqrt{\left(20 - \frac{12}{x} \right)}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 2

Question:

A particle P is moving along the x -axis. At time t seconds, the displacement of P from the origin O is x m and the velocity of P is $4e^{0.5t}$ m s⁻¹ in the direction Ox . When $t = 0$, P is at O . Find

- x in terms of t ,
- the acceleration of P when $t = \ln 9$.

Solution:

$$\text{a } v = \frac{dx}{dt} = 4e^{0.5t}$$

$$x = \int 4e^{0.5t} dt = 8e^{0.5t} + A$$

When $t = 0$, $x = 0$

$$0 = 8 + A \Rightarrow A = -8$$

$$x = 8e^{0.5t} - 8 = 8(e^{0.5t} - 1)$$

$$\text{b } a = \frac{dv}{dt} = 2e^{0.5t}$$

When $t = \ln 9$

$$a = 2e^{0.5 \ln 9} = 2e^{\ln 3} = 2 \times 3 = 6$$

The acceleration of P when $t = \ln 9$ is 6 m s^{-2} .

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 3

Question:

A particle is moving along the x -axis. At time $t = 0$, P is passing through the origin O with velocity 8 m s^{-1} in the direction of x increasing. When P is $x \text{ m}$ from O , its

acceleration is $\left(3 + \frac{1}{4}x\right) \text{ m s}^{-2}$ in the direction of x decreasing.

Find the positive value of x for which P is instantaneously at rest.

Solution:

$$a = -\left(3 + \frac{1}{4}x\right)$$

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -3 - \frac{1}{4}x$$

$$\frac{1}{2}v^2 = \int\left(-3 - \frac{1}{4}x\right)dx = -3x - \frac{1}{8}x^2 + A$$

$$\text{At } x = 0, v = 8$$

$$32 = -0 - 0 + A \Rightarrow A = 32$$

$$\frac{1}{2}v^2 = 32 - 3x - \frac{1}{8}x^2$$

$$\text{When } v = 0$$

$$0 = 32 - 3x - \frac{1}{8}x^2$$

$$x^2 + 24x - 256 = 0$$

$$(x + 32)(x - 8) = 0$$

$$\text{As } x > 0$$

$$x = 8$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 4

Question:

A particle P is moving on the x -axis. When P is a distance x metres from the origin O , its acceleration is of magnitude $\frac{15}{4x^2} \text{ m s}^{-2}$ in the direction OP . Initially P is at the point where $x = 5$ and is moving toward O with speed 6 m s^{-1} . Find the value of x where P first comes to rest.

Solution:

$$a = \frac{15}{4x^2} = \frac{15}{4} x^{-2}$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = \frac{15}{4} x^{-2}$$

$$\frac{1}{2} v^2 = \int \frac{15}{4} x^{-2} dx = -\frac{15}{4} x^{-1} + A$$

$$\frac{1}{2} v^2 = A - \frac{15}{4x}$$

At $x = 5, v = -6$

$$18 = A - \frac{15}{20} \Rightarrow A = 18\frac{3}{4} = \frac{75}{4}$$

$$\frac{1}{2} v^2 = \frac{75}{4} - \frac{15}{4x} = \frac{15}{4} \left(5 - \frac{1}{x} \right)$$

When $v = 0$

$$5 - \frac{1}{x} = 0 \Rightarrow x = \frac{1}{5}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 5

Question:

A particle P moves along the x -axis in the direction x increasing. At time t seconds, the velocity of P is v m s⁻¹ and its acceleration is $20te^{-t^2}$ m s⁻². When $t = 0$ the speed of P is 8 m s⁻¹. Find

- v in terms of t ,
- the limiting velocity of P .

Solution:

a $a = \frac{dv}{dt} = 20te^{-t^2}$

$$v = \int 20te^{-t^2} dt = -10e^{-t^2} + A$$

When $t = 0, v = 8$

$$8 = -10 + A \Rightarrow A = 18$$

$$v = 18 - 10e^{-t^2}$$

b As $t \rightarrow \infty, e^{-t^2} \rightarrow 0$ and $v \rightarrow 18$

The limiting velocity of P is 18 m s⁻¹.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 6

Question:

A particle P moves along a straight line. Initially P is at rest at a point O on the line.

A time t seconds, where $t \geq 0$, the acceleration of P is $\frac{18}{(2t+3)^3} \text{ m s}^{-2}$ directed away

from O .

Find the value of t for which the speed of P is 0.48 ms^{-1} .

Solution:

$$a = \frac{dv}{dt} = \frac{18}{(2t+3)^3} = 18(2t+3)^{-3}$$

$$\begin{aligned} v &= \int 18(2t+3)^{-3} dt = \frac{18}{-2 \times 2} (2t+3)^{-2} + A \\ &= A - \frac{9}{2(2t+3)^2} \end{aligned}$$

When $t = 0, v = 0$

$$0 = A - \frac{9}{2 \times 3^2} \Rightarrow A = \frac{1}{2}$$

$$v = \frac{1}{2} - \frac{9}{2(2t+3)^2}$$

When $v = 0.48$

$$0.48 = \frac{1}{2} - \frac{9}{2(2t+3)^2} \Rightarrow \frac{9}{2(2t+3)^2} = 0.02$$

$$(2t+3)^2 = \frac{9}{2 \times 0.02} = 225$$

$$t \geq 0$$

As $2t+3 = \sqrt{225} = 15$

$$t = \frac{15-3}{2} = 6$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 7

Question:

A particle P is moving along the x -axis. At time t seconds, the velocity of P is v m s⁻¹ and the acceleration of P is $(3-x)$ m s⁻² in the direction x increasing. Initially P is at the origin O and is moving with speed 4 m s⁻¹ in the direction x increasing. Find

- v^2 in terms of x ,
- the maximum value of v .

Solution:

a $a = 3 - x$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 3 - x$$

$$\frac{1}{2} v^2 = \int (3 - x) dx = 3x - \frac{x^2}{2} + A$$

$$v^2 = B + 6x - x^2, \text{ where } B = 2A$$

At $x = 0, v = 4$

$$16 = B + 0 - 0 \Rightarrow B = 16$$

$$v^2 = 16 + 6x - x^2$$

b $v^2 = 16 + 6x - x^2 = 25 - 9 + 6x - x^2$
 $= 25 - (x - 3)^2$

As $(x - 3)^2 \geq 0, v^2 \leq 25$

The greatest value of v is 5.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 8

Question:

A particle P is moving along the x -axis. At time $t = 0$, P passes through the origin O . After t seconds the speed of P is v m s⁻¹, $OP = x$ metres and the acceleration of P is $\frac{x^2(5-x)}{2}$ m s⁻² in the direction x increasing. At $x = 10$, P is instantaneously at rest.

Find

- an expression for v^2 in terms of x ,
- the speed of P when $t = 0$.

Solution:

$$\text{a } a = \frac{d}{dx} \left(\frac{1}{2} v^2 \right) = \frac{x^2(5-x)}{2} = \frac{5x^2}{2} - \frac{x^3}{2}$$

$$\frac{1}{2} v^2 = \int \left(\frac{5x^2}{2} - \frac{x^3}{2} \right) dx = \frac{5x^3}{6} - \frac{x^4}{8} + A$$

$$v^2 = \frac{5x^3}{3} - \frac{x^4}{4} + B, \text{ where } B = 2A$$

At $x = 10$, $v = 0$

$$0 = \frac{5000}{3} - \frac{10\,000}{4} + B \Rightarrow B = \frac{2500}{3}$$

$$v^2 = \frac{5x^3}{3} - \frac{x^4}{4} + \frac{2500}{3}$$

- When $t = 0$, $x = 0$

$$v^2 = \frac{2500}{3} \Rightarrow v = (\pm) \frac{50}{\sqrt{3}} = (\pm) \frac{50\sqrt{3}}{3}$$

The speed of P when $t = 0$ is $\frac{50\sqrt{3}}{3}$ m s⁻¹.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 9

Question:

A particle P moves away from the origin along the positive x -axis. At time t seconds, the acceleration of P is $\frac{20}{5x+2}$ m s⁻², where $OP = x$ m, directed away from O . Given

that the speed of P is 3 m s⁻¹ at $x = 0$, find, giving your answers to 3 significant figures,

- the speed of P at $x = 12$,
- the value of x when the speed of P is 5 m s⁻¹.

Solution:

$$\text{a } a = \frac{d}{dx} \left(\frac{1}{2} v^2 \right) = \frac{20}{5x+2}$$

$$\frac{1}{2} v^2 = \int \frac{20}{5x+2} dx = 4 \ln(5x+2) + A$$

$$v^2 = 8 \ln(5x+2) + B, \text{ where } B = 2A$$

$$\text{At } x = 0, v = 3$$

$$9 = 8 \ln 2 + B \Rightarrow B = 9 - 8 \ln 2$$

$$v^2 = 8 \ln(5x+2) - 8 \ln 2 + 9 = 8 \ln \left(\frac{5x+2}{2} \right) + 9$$

$$\text{At } x = 12$$

$$v^2 = 8 \ln 31 + 9 = 36.471 \dots$$

$$v = \sqrt{36.471 \dots} = 6.039 \dots$$

The speed of P at $x = 12$ is 6.04 m s⁻¹ (3 s.f.).

- When $v = 5$

$$25 = 8 \ln \left(\frac{5x+2}{2} \right) + 9$$

$$\ln \left(\frac{5x+2}{2} \right) = \frac{25-9}{8} = 2$$

$$\frac{5x+2}{2} = e^2$$

$$x = \frac{2e^2 - 2}{5} = 2.56 \text{ (3 s.f.)}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 10

Question:

A car moves along a horizontal straight road. At time t seconds the acceleration of the car is $\frac{100}{(2t+5)^2}$ m s⁻² in the direction of motion of the car. When $t = 0$, the car is at rest. Find

- an expression for v in terms of t ,
- the distance moved by the car in the first 10 seconds of its motion.

Solution:

$$\text{a } a = \frac{dv}{dt} = \frac{100}{(2t+5)^2} = 100(2t+5)^{-2}$$

$$v = \int 100(2t+5)^{-2} dt = \frac{100}{2 \times -1} (2t+5)^{-1} + A$$

$$= A - \frac{50}{2t+5}$$

$$\text{When } t = 0, v = 0$$

$$0 = A - \frac{50}{5} \Rightarrow A = 10$$

$$v = 10 - \frac{50}{2t+5}$$

$$\text{b } v = \frac{dx}{dt} = 10 - \frac{50}{2t+5}$$

$$x = \int \left(10 - \frac{50}{2t+5} \right) dt = 10t - 25 \ln(2t+5) + B$$

$$\text{When } t = 0, x = 0$$

$$0 = -25 \ln 5 + B \Rightarrow B = 25 \ln 5$$

$$x = 10t - 25 \ln(2t+5) + 25 \ln 5$$

$$\text{When } t = 10$$

$$x = 100 - 25 \ln 25 + 25 \ln 5 = 100 - 25 \ln \frac{25}{5} = 100 - 25 \ln 5$$

The distance moved by the car in the first 10 seconds of its motion is $(100 - 25 \ln 5)$ m (≈ 59.8 m).

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 11

Question:

A particle P is moving in a straight line with acceleration $\cos^2 t \text{ m s}^{-2}$ at time t seconds. The particle is initially at rest at a point O .

a Find the speed of P when $t = \pi$.

b Show that the distance of P from O when $t = \frac{\pi}{4}$ is $\frac{1}{64}(\pi^2 + 8)\text{m}$.

Solution:

$$\text{a } a = \frac{dv}{dt} = \cos^2 t = \frac{1}{2} + \frac{1}{2} \cos 2t$$

$$v = \int \left(\frac{1}{2} + \frac{1}{2} \cos 2t \right) dt = \frac{1}{2}t + \frac{1}{4} \sin 2t + A$$

When $t = 0, v = 0$

$$0 = 0 + 0 + A \Rightarrow A = 0$$

$$v = \frac{1}{2}t + \frac{1}{4} \sin 2t$$

When $t = \pi$

$$v = \frac{\pi}{2} + \frac{1}{4} \sin 2\pi = \frac{\pi}{2} + 0 = \frac{\pi}{2}$$

The speed of P when $t = \pi$ is $\frac{\pi}{2} \text{ m s}^{-1}$.

b The distance of P from O when $t = \frac{\pi}{4}$ is given by

$$x = \int_0^{\frac{\pi}{4}} \left(\frac{1}{2}t + \frac{1}{4} \sin 2t \right) dt = \left[\frac{1}{4}t^2 - \frac{1}{8} \cos 2t \right]_0^{\frac{\pi}{4}}$$

$$= \left(\frac{\pi^2}{64} - \frac{1}{8} \cos \frac{\pi}{2} \right) - \left(0 - \frac{1}{8} \right)$$

$$= \frac{\pi^2}{64} + \frac{1}{8} = \frac{1}{64}(\pi^2 + 8)$$

The distance of P from O when $t = \frac{\pi}{4}$ is $\frac{1}{64}(\pi^2 + 8)\text{m}$, as required.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 12

Question:

A particle P is moving along the x -axis. At time t seconds, the velocity of P is $v \text{ m s}^{-1}$ in the direction of x increasing, where

$$v = \begin{cases} \frac{1}{2}t^2, & 0 \leq t \leq 4 \\ 8e^{4-t}, & t > 4 \end{cases}$$

When $t = 0$, P is at the origin O . Find

- the acceleration of P when $t = 2.5$,
- the acceleration of P when $t = 5$,
- the distance of P from O when $t = 6$.

Solution:

a When $t = 2.5$, $v = \frac{1}{2}t^2$

$$a = \frac{dv}{dt} = t$$

When $t = 2.5$, $a = 2.5$

The acceleration of P when $t = 2.5$ is 2.5 m s^{-2} in the direction of x increasing.

b When $t = 5$, $v = 8e^{4-t}$

$$a = \frac{dv}{dt} = -8e^{4-t}$$

When $t = 5$, $a = -8e^{4-5} = -8e^{-1}$

The acceleration of P when $t = 5$ is $8e^{-1} \text{ m s}^{-2}$ in the direction of x decreasing.

- c The distance of P from O when $t = 6$ is given by

$$\begin{aligned} x &= \int_0^4 \frac{1}{2}t^2 dt + \int_4^6 8e^{4-t} dt \\ &= \left[\frac{t^3}{6} \right]_0^4 + \left[-8e^{4-t} \right]_4^6 = \frac{64}{6} - 8e^{-2} + 8 \\ &= \frac{56}{3} - 8e^{-2} \end{aligned}$$

The distance of P from O when $t = 6$ is $\left(\frac{56}{3} - 8e^{-2} \right) \text{ m} \approx 17.6 \text{ m}$ (3 s.f.).

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 13

Question:

A particle P is moving along the x -axis. When $t = 0$, P is passing through O with velocity 3 m s^{-1} in the direction of x increasing. When $0 \leq x \leq 4$ the acceleration is of magnitude $\left(4 + \frac{1}{2}x\right) \text{ m s}^{-2}$ in the direction of x increasing. At $x = 4$, the acceleration of P changes. For $x > 4$, the magnitude of the acceleration remains $\left(4 + \frac{1}{2}x\right) \text{ m s}^{-2}$ but it is now in the direction of x decreasing.

- Find the speed of P at $x = 4$.
- Find the positive value of x for which P is instantaneously at rest. Give your answer to 2 significant figures.

Solution:

$$\mathbf{a} \quad a = \frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 4 + \frac{1}{2} x$$

$$\frac{1}{2} v^2 = 4x + \frac{x^2}{4} + A$$

$$v^2 = 8x + \frac{x^2}{2} + B, \text{ where } B = 2A$$

$$\text{At } x = 0, v = 3$$

$$9 = 0 + 0 + B \Rightarrow B = 9$$

$$v^2 = 8x + \frac{x^2}{2} + 9$$

$$\text{At } x = 4$$

$$v^2 = 32 + 8 + 9 = 49 \Rightarrow v = 7$$

The speed of P at $x = 4$ is 7 m s^{-1} .

$$\mathbf{b} \quad a = \frac{d}{dx} \left(\frac{1}{2} v^2 \right) = -4 - \frac{1}{2} x$$

$$\frac{1}{2} v^2 = C - 4x - \frac{x^2}{4}$$

$$v^2 = D - 8x - \frac{x^2}{2}, \text{ where } D = 2C$$

$$\text{At } x = 4, v = 7$$

$$49 = D - 32 - 8 \Rightarrow D = 89$$

$$v^2 = 89 - 8x - \frac{x^2}{2}$$

$$\text{When } v = 0$$

$$x^2 + 16x = 178 \Rightarrow x^2 + 16x + 64 = 242$$

$$(x + 8)^2 = 242 \Rightarrow x = 11\sqrt{2} - 8, \text{ as } x > 0$$

$$x = 7.6 \text{ (2 s.f.)}$$

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 14

Question:

A particle P is moving along the x -axis. At time t seconds, P has velocity v m s⁻¹ in

the direction x increasing and an acceleration of magnitude $\frac{2t+3}{t+1}$ m s⁻² in the

direction x increasing. When $t = 0$, P is at rest at the origin O . Find

- v in terms of t ,
- the distance of P from O when $t = 2$.

Solution:

$$\text{a } \alpha = \frac{dv}{dt} = \frac{2t+3}{t+1} = 2 + \frac{1}{t+1}$$

$$v = 2t + \ln(t+1) + A$$

$$\text{When } t = 0, v = 0$$

$$0 = 0 + A \Rightarrow A = 0$$

$$v = 2t + \ln(t+1)$$

- The distance of P from O when $t = 2$ is given by

$$x = \int_0^2 (2t + \ln(t+1)) dt$$

Using integration by parts

$$\int \ln(t+1) dt = \int 1 \ln(t+1) dt = t \ln(t+1) - \int \frac{t}{t+1} dt$$

$$= t \ln(t+1) - \int \left(1 - \frac{1}{t+1} \right) dt = t \ln(t+1) - t + \ln(t+1)$$

$$= (t+1) \ln(t+1) - t (+C)$$

$$\text{Hence } x = \left[t^2 - t + (t+1) \ln(t+1) \right]_0^2 = 2 + 3 \ln 3$$

The distance of P from O when $t = 2$ is $(2 + 3 \ln 3)$ m.

Solutionbank M3

Edexcel AS and A Level Modular Mathematics

Further kinematics

Exercise C, Question 15

Question:

A particle P is moving along the x -axis. At time t seconds P is x m from O , has velocity v m s⁻¹ and acceleration of magnitude $(4x+6)$ m s⁻² in the direction of x increasing. When $t=0$, P is passing through O with velocity 3 m s⁻¹ in the direction of x increasing. Find

- v in terms of x ,
- x in terms of t .

Solution:

$$\text{a } a = \frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 4x + 6$$

$$\frac{1}{2} v^2 = 2x^2 + 6x + A$$

$$v^2 = 4x^2 + 12x + B, \text{ where } B = 2A$$

$$\text{At } x = 0, v = 3$$

$$9 = 0 + 0 + B \Rightarrow B = 9$$

$$v^2 = 4x^2 + 12x + 9 = (2x + 3)^2$$

As v is increasing as x increases

$$v = 2x + 3$$

$$\text{b } v = \frac{dx}{dt} = 2x + 3$$

Separating the variables and integrating

$$\int \frac{1}{2x+3} dx = \int 1 dt$$

$$\frac{1}{2} \ln(2x+3) = t + C$$

$$\ln(2x+3) = 2t + 2C$$

$$2x+3 = e^{2t+2C} = D e^{2t}, \text{ where } D = e^{2C}$$

When $t = 0, x = 0$

$$3 = D e^0 \Rightarrow D = 3$$

$$2x+3 = 3e^{2t}$$

$$x = \frac{3}{2} (e^{2t} - 1)$$