

GCE Examinations

Pure Mathematics

Module P5

Advanced Subsidiary / Advanced Level

Paper H

Time: 1 hour 30 minutes

Instructions and Information

Candidates may use any calculator except those with a facility for symbolic algebra and/or calculus.

Full marks may be obtained for answers to ALL questions.

Mathematical and statistical formulae and tables are available.

This paper has 7 questions.

Advice to Candidates

You must show sufficient working to make your methods clear to an examiner.
Answers without working will gain no credit.



Written by Rosemary Smith & Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

1. A curve has the equation

$$2x^2 + y^2 = 4.$$

Find the radius of curvature of the curve at the point $(1, -\sqrt{2})$.

(8 marks)

2. (a) Using the definition of $\cosh x$ in terms of exponential functions show that $\cosh x$ is an even function.

(2 marks)

- (b) Given that $x > 0$ and $y > 0$, solve the simultaneous equations

$$\ln(xy) = \operatorname{arcosh}\left(\frac{5}{3}\right),$$

$$\cosh(3x - y) = 1.$$

(6 marks)

3. Find

$$\int \frac{1}{13 \cosh x - 5 \sinh x} dx.$$

(8 marks)

4. (a) Given that $y = \arcsin(2x - 1)$, prove that

$$\frac{dy}{dx} = \frac{1}{\sqrt{x - x^2}}.$$

(4 marks)

The tangent to the curve $y = \arcsin(2x - 1)$ at the point where $x = \frac{3}{4}$ meets the y -axis at A .

- (b) Find the exact value of the y -coordinate of A .

(5 marks)

5. The point $P(at^2, 2at)$, $t \neq 0$, lies on the parabola C with equation $y^2 = 4ax$.

(a) Show that an equation of the tangent to C at P is

$$yt = x + at^2. \quad (4 \text{ marks})$$

The tangent to C at P meets the x -axis at Q and the y -axis at R .

M is the mid-point of QR .

(b) Find the coordinates of M . (3 marks)

Given that OM is perpendicular to OP , where O is the origin,

(c) show that $t^2 = 2$. (4 marks)

6.
$$I_n = \int \frac{\cos n\theta}{\sin \theta} d\theta, \quad n \in \mathbb{N}.$$

(a) By considering $I_n - I_{n-2}$, or otherwise, show that

$$I_n = \frac{2 \cos(n-1)\theta}{n-1} + I_{n-2}. \quad (5 \text{ marks})$$

(b) Hence evaluate

$$\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{\cos 5\theta}{\sin \theta} d\theta,$$

leaving your answer in terms of natural logarithms. (8 marks)

Turn over

7. The ellipse C has equation $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where a and b are positive constants and $a > b$.

The coordinates of the foci of C are $(\pm\sqrt{3}, 0)$, and the equations of its directrices are $x = \pm \frac{4}{\sqrt{3}}$.

- (a) Find the value of a and the value of b . **(4 marks)**

The ellipse is rotated completely about the x -axis.

- (b) Show that the area of the surface of revolution generated is given by

$$A = \frac{\pi}{2} \int_{-2}^2 \sqrt{16 - 3x^2} \, dx. \quad \text{(6 marks)}$$

- (c) Use integration to show that

$$A = \frac{8}{9} \pi^2 \sqrt{3} + 2\pi. \quad \text{(8 marks)}$$

END