

ADVANCED SUBSIDIARY GCE

CHEMISTRY

Practical Examination 1 (Part B – Practical Test)

MONDAY 12 MAY 2008

2813/03/TEST

Afternoon

Time: 1 hour 30 minutes

Candidates answer on the question paper.

Additional materials:

Candidate's Plan (Part A of the Practical Examination)

Scientific Calculator

Data Sheet for Chemistry (Inserted)



Candidate
Name

Centre
Number

--	--	--	--	--

Candidate
Number

--	--	--	--

INSTRUCTIONS TO CANDIDATES

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer **all** the questions.
- Write your answers in the spaces provided on the question paper.
- Read the instructions and questions carefully.
- Do not write in the bar code. Do not write in the grey area between the pages.

INFORMATION FOR CANDIDATES

- In this part of the Practical Test, you will be assessed on the Experimental and Investigative Skills:
Skill I Implementing
Skill A Analysing evidence and drawing conclusions
Skill E Evaluating evidence and procedures
- You may use a scientific calculator.
- You are advised to show all the steps in any calculations.
- You may refer to your Plan produced for Part A.
- You will be awarded marks for the quality of written communication where this is indicated.
- A copy of the *Data Sheet for Chemistry* is provided as an insert with this question paper.

FOR EXAMINER'S USE		
Qu.	Max.	Mark
Planning	16	
Implementing & Analysing	30	
Evaluating	14	
TOTAL	60	

This document consists of **10** printed pages, **2** blank pages and a *Data Sheet for Chemistry*.

Answer **all** the parts.

Introduction

In this examination, you will investigate the reaction between sulphamic acid, H_3NSO_3 , and potassium hydroxide by carrying out a titration.

Three chemicals are provided.

- Solution **Y** is aqueous potassium hydroxide, KOH .
- Solid **Z** is sulphamic acid, H_3NSO_3 .
- Phenolphthalein solution is the indicator.

Irritant



Irritant



Flammable



Part 1 Titration of aqueous KOH with aqueous H_3NSO_3 Skill I (Implementing)

[16 marks]

Record all your readings in the spaces on page 3.

Weigh the bottle containing **Z**, with the lid.

Tip all of **Z** into a 250 cm^3 beaker.

Weigh the empty bottle and lid.

Calculate the mass of **Z** used.

In the beaker, dissolve **Z** in about 100 cm^3 of distilled (or de-ionised) water.

When the solid has dissolved, transfer the solution into a 250 cm^3 volumetric flask.

Use distilled water to rinse any remaining solution from the beaker into the flask.

Make the solution up to 250 cm^3 using distilled water.

Invert the volumetric flask several times before use, to mix the solution thoroughly.

Using a pipette and filler, transfer 25.0 cm^3 of this solution of **Z** into a conical flask.

Add five drops of phenolphthalein indicator.

Fill the burette with solution **Y**.

Record all burette readings to 0.05 cm^3 .

The end-point of the titration is when the indicator changes from colourless to pale pink.

Carry out a trial titration.

Repeat the titration until you obtain **two** consistent accurate results.

In each case, use 25.0 cm^3 of your solution of **Z**.

Calculate your mean titre.

Readings and calculations

Use the spaces below to record **all** your readings.

Weighings

mass of **Z** used = g

Titration data (tabulated)**Summary**

25.0 cm³ of the solution of **Z** required a mean titre of cm³ of **Y**.

Show the readings you used to obtain this value of the volume of **Y** by putting a tick (✓) under these readings.

Safety

A bottle of **solid** potassium hydroxide is labelled as **Corrosive**.

Suggest why the bottle containing **Y**, aqueous potassium hydroxide, is labelled **Irritant**.

.....

.....

.....

.....

Part 2 **Deducing the equation for the reaction of KOH with H_3NSO_3**
Skill A (Analysing)

[14 marks]

Use the *Data Sheet for Chemistry* supplied for any data you require.
Your working must be shown clearly.
Give your answers to **three** significant figures for parts **(a) – (e)**.

- (a)** Solution **Y** contains 5.50 g dm^{-3} of **impure** KOH.
The KOH used to make up solution **Y** was 86.0% pure by mass.

Calculate the mass of pure KOH in 1.00 dm^3 of solution **Y**.

mass of pure KOH = g

- (b)** Calculate the concentration, in mol dm^{-3} , of pure KOH in solution **Y**.

concentration = mol dm^{-3}

- (c)** Calculate the number of moles of KOH in the mean titre.

answer = mol

- (d) Calculate the number of moles of sulphamic acid, H_3NSO_3 , **Z**, that you weighed out.

answer = mol

- (e) Calculate the number of moles of H_3NSO_3 , **Z**, used in each titration.

answer = mol

- (f) Use your answers to (c) and (e) to calculate how many moles of KOH react with 1 mol of H_3NSO_3 .




Show your working and give your answer to the nearest whole number.

answer = mol KOH

- (g) Three possible reactions of KOH with H_3NSO_3 , are represented by the equations below.

- (i) Complete and balance the second and third equations.

State symbols are **not** required.



- (ii) Write a tick (✓) next to the equation which matches most closely to your answer to (f).

Explain your choice.

.....

- (h) Sulphamic acid is a strong acid.

Deduce the number of moles of hydrogen ions, H^+ , present in an aqueous solution containing 1 mole of H_3NSO_3 .


answer = mol

Information


A student carried out a different experiment to determine the number of moles of H^+ ions in an aqueous solution containing 1 mole of aqueous sulphamic acid, **Z**.

He added excess magnesium to a solution of **Z**, and measured the volume of hydrogen produced in a 100 cm^3 gas syringe.

The apparatus used is shown below.



- (a) The equations below represent three possible reactions of magnesium with aqueous sulphamic acid, **Z**.



In a preliminary experiment, which was carried out at room temperature and pressure (r.t.p.), the student used 0.97 g of **Z**, dissolved in water.

Carry out a calculation to find out whether this was a suitable mass of **Z** to use with a 100 cm^3 gas syringe. The volume of 1 mole of any gas at r.t.p. = $24\,000\text{ cm}^3$.

Explain your working.

- (b) State why it is important that an **excess** of magnesium is used during the reaction.

.....

.....

..... [1]

- (c) Suggest **three** sources of error in the student's experiment.

Explain how **each** source of error could be reduced or minimised.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... [6]

- (d) The student carried out the experiment twice.

His readings are shown in the table below.

mass of sulphamic acid, Z , used/g	0.64	0.45
volume of gas collected/cm ³	80	50

- (i) Carry out a calculation(s) to find out whether the student should repeat the experiment again.

[2]

- (ii) Which experiment gives more **reliable** results, your own (involving the titration) or the student's?

Justify your answer.

.....

.....

.....

.....

.....

..... [2]

END OF QUESTION PAPER

10
BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

11
BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

PLEASE DO NOT WRITE ON THIS PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© OCR 2008