

Mark Scheme 1	Matching the syllabus written by EDEXCEL Curriculum 2004+
Calculators Allowed <i>Where appropriate leave your answers to 3 s.f.</i>	
© ZigZag Education 2004	Core Mathematics – C4

1. Use identity $\cos^2 A + \sin^2 A = 1 \Rightarrow \cos^2 A = 1 - \sin^2 A$ M1
 And substitute into $\cos 2x = \cos^2 x - \sin^2 x$
 $\cos 2x = 1 - 2\sin^2 x$
 $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$
 Separating variables: $\int 1 dy = \int (\sin^2 x) dx$ M1
 Hence $y = \frac{1}{2} \int (1 - \cos 2x) dx$
 $= \frac{1}{2} \left(x - \frac{1}{2} \sin 2x \right) + c$ A1A1
 Substitute $x = 0.1$, $y = 0.2$ to obtain $c = 0.19966\dots = 0.200$ (3 s.f.) M1
 i.e. $y = \frac{1}{2} \left(x - \frac{1}{2} \sin 2x \right) + 0.200$ A1 (6)

Note: alternative solution using integration by parts: $y = \int \sin x \sin x dx$

2. a) Expand binomial $(1+x)^n = 1 + nx + \frac{n(n-1)x^2}{1 \times 2} + \frac{n(n-1)(n-2)x^3}{1 \times 2 \times 3} + \dots$ M1
 $(1-x)^{\frac{1}{2}} = 1 + \frac{1}{2}(-x) + \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(\frac{1}{2}\right)(-x)^2 + \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(\frac{1}{6}\right)(-x)^3$ M1
 $= 1 - \frac{1}{2}x - \frac{1}{8}x^2 - \frac{1}{16}x^3$ A1A1(4)

b) Substitute $x = 10^{-2}$ into expansion from a) to obtain:

$$\left(1 - \frac{1}{100}\right)^{\frac{1}{2}} = 0.994987\dots$$

M1 ft

$$\frac{1}{10} \sqrt{99} = 0.994987\dots$$

$$\frac{3}{10} \sqrt{11} = 0.994987\dots$$

M1

$$\text{Hence } \sqrt{11} = 3.3166 \text{ (5 s.f.)}$$

A1 (3)

3. a) Attempt integration by parts M1
 with $u = x$ $v' = e^x$ M1
 $\int (xe^x) dx = xe^x - \int (1 \cdot e^x) dx$ A1A1
 $= xe^x - e^x + c$ A1 (5)
- b) Attempt integration by parts twice M1
 $\int (x^2 e^x) dx = x^2 e^x - \int (2xe^x) dx$ A1A1
 $= x^2 e^x - 2(xe^x - e^x)$ substitute answer from a) M1
 $= x^2 e^x - 2xe^x + 2e^x + c$ A1 (5)

4. a) Use scalar product formula $\mathbf{a} \cdot \mathbf{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$ M1
 Vectors are perpendicular $\therefore \mathbf{a} \cdot \mathbf{b} = 0$ M1
 Hence $-1 \times 1 + 1 \times c + 1 \times 1 = 0$ M1

Answer $c = 0$ A1 (4)

b) Form 2 simultaneous equations:
 $1 + 3\mu = 1 + \lambda$ M1
 $2 + 2\mu = \lambda$ M1
Solve simultaneous equations to find $\mu = 2$ or $\lambda = 6$ M1
Substitute one of these back into the appropriate line equation to find
position vector of intersection point: $7\mathbf{i} + 6\mathbf{j} + 7\mathbf{k}$ A1 (4)

c) Use formula $\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3 = |\mathbf{a}||\mathbf{b}|\cos\theta$ with $3\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ and $\mathbf{i} + \mathbf{j} + \mathbf{k}$ M1
i.e. $\cos\theta = \frac{(3 \times 1) + (2 \times 1) + (3 \times 1)}{\sqrt{3^2 + 2^2 + 3^2} \sqrt{1^2 + 1^2 + 1^2}}$ M1
 $\therefore \theta = \cos^{-1}\left(\frac{8}{\sqrt{66}}\right) = 10.024\dots = 10.0^\circ$ (3 s.f.) A1 (3)

5. a) Write as $f(x) = \frac{A}{x-1} + \frac{B}{x+2}$ M1
Therefore $x + 1 = (x + 2)A + (x - 1)B$ M1
Substitute $x = 1$, or similar method, (to find $A = \frac{2}{3}$) M1
Substitute $x = -2$, or similar method, (to find $B = \frac{1}{3}$) M1
Hence $f(x) = \frac{2}{3(x-1)} + \frac{1}{3(x+2)}$ A1A1(6)

b) $\frac{df}{dx} = -\frac{2}{3(x-1)^2} - \frac{1}{3(x+2)^2}$ A1A1
Substituting $x = 2$, $\frac{df(2)}{dx} = -\frac{2}{3} - \frac{1}{48} = -\frac{11}{16}$ M1A1(4)

c) $\int_4^5 \frac{2}{3(x-1)} + \frac{1}{3(x+2)} dx$
 $= \left[\frac{2}{3} \ln(x-1) + \frac{1}{3} \ln(x+2) \right]_4^5$ M1M1
 $= \frac{2}{3} \ln 4 + \frac{1}{3} \ln 7 - \frac{2}{3} \ln 3 - \frac{1}{3} \ln 6 = \frac{2}{3} \ln\left(\frac{4}{3}\right) + \frac{1}{3} \ln\left(\frac{7}{6}\right)$ or $= \frac{1}{3} \ln\left(\frac{56}{27}\right)$ M1A1(4)

6. a) Substitute $x = \frac{1}{2}$ into $x^2 + y^2 + 2x + 4y + 1 = 0$ M1
Simplify to quadratic $y^2 + 4y + \frac{9}{4} = 0$ M1
Use quadratic formula to solve for y
Hence coordinates $(0.5, -0.67712\dots)$ and $(0.5, -3.3228\dots)$
 $= (0.5, -0.68)$ and $(0.5, -3.32)$ (3 s.f.) A1 (3)

b) Differentiating implicitly,
 $2x + 2yy' + 2 + 4y' = 0$ A1A1A1A1
 $y'(2y + 4) = -2x - 2$ M1
 $\therefore y' = \frac{-2x - 2}{2y + 4} = -\frac{(x+1)}{y+2}$ A1 (6)

c) Substituting coordinates $(0.5, -0.67\dots)$,

$$y' = \frac{-2x-2}{2y+4} = -\frac{(x+1)}{y+2} = -\frac{3\sqrt{7}}{7} = -1.1338\dots = -1.13 \text{ (3 s.f.)} \quad \text{M1A1}$$

Substituting coordinates (0.5, -3.3...),

$$y' = \frac{-2x-2}{2y+4} = -\frac{(x+1)}{y+2} = \frac{3\sqrt{7}}{7} = 1.1338\dots = 1.13 \text{ (3 s.f.)} \quad \text{M1A1(4)}$$

- d) $y = 1.1\dots x + c$
 Substitute (0.5, -3.3...) to find c M1
 $c = -3.8898\dots = 3.89 \text{ (3 s.f.)}$
 and hence $y = 1.13x - 3.89 \text{ (3s.f.)}$ A1 (2)

7. $V = \pi \int y^2 dx$ M1

$$= \pi \int_0^\pi x^2 \sin x dx \quad \text{limits M1A1}$$

$$\int v \frac{du}{dx} dx = uv - \int u \frac{dv}{dx} dx$$

$$\frac{du}{dx} = \sin x \quad v = x^2 \quad \text{M1}$$

$$u = -\cos x \quad \frac{dv}{dx} = 2x \quad \text{M1}$$

$$V = -\cos x \times x^2 - \int -\cos x(2x) dx \quad \text{A1}$$

$$I = \int 2x \cos x dx \quad \frac{du}{dx} = \cos x \quad v = 2x$$

$$u = \sin x \quad \frac{dv}{dx} = 2 \quad \text{M1}$$

$$I = 2x \sin x - 2 \int \sin x dx \quad \text{M1}$$

$$= 2x \sin x + 2 \cos x \quad \text{A1}$$

$$\text{Therefore } V = \pi \left[-x^2 \cos x + 2x \sin x + 2 \cos x \right]_0^\pi \quad \text{M1}$$

$$= \pi \left[-\pi^2 \cos \pi + 2\pi \sin \pi + 2 \cos \pi - (2 \cos 0) \right] \quad \text{M1}$$


$$= \pi \left[\pi^2 - 2 - 2 \right]$$

$$= \pi \left[\pi^2 - 4 \right] \text{ cm}^3 \text{ or equivalent} \quad \text{A1 (12)}$$

(75)

Mark Scheme 2	Matching the syllabus written by EDEXCEL Curriculum 2004+
Calculators Allowed <i>Where appropriate leave your answers to 3 s.f.</i>	
© ZigZag Education 2004	Core Mathematics – C4

1. a) Curve sketch which cuts the y-axis at $y = 3$ A1A1(2)



- b) The area is $\int_0^{\ln 10} 3e^x dx = [3e^x]_0^{\ln 10} = 27$ square units M1A1M1A1
(4)

2. a) $\int_1^4 \frac{1}{1+2x} dx = \int_1^4 \frac{1}{y} \frac{dx}{dy} dy$ M1
 $y = 1 + 2x, \frac{dy}{dx} = 2, \frac{dx}{dy} = \frac{1}{2}$ M1
 $\int_1^4 \frac{1}{y} \frac{dx}{dy} dy = \left[\frac{1}{2} \ln|1+2x| \right]_1^4 = \frac{\ln 9}{2} - \frac{\ln 3}{2} = \frac{\ln 3}{2}$ A1 (3)

- b) Using the trapezium rule correctly, with $n = 3$. i.e. $\frac{h}{2}[y_0 + 2y_1 + 2y_2 + y_3]$ M1
Using $f(1), f(2), f(3)$ or $f(4)$ M1
 $h = 1; [x_0 = \frac{1}{3}; x_1 = \frac{2}{5}; x_2 = \frac{3}{7}; x_3 = \frac{4}{9}]$ A1[A1]
Estimated area = $\frac{1}{2} \left[\frac{1}{3} + \frac{4}{9} + 2 \left(\frac{2}{5} + \frac{3}{7} \right) \right] = 1 \frac{137}{630}$ (= 1.2174... = 1.22 square units to 3 s.f.) A1 (5)

3. a) Using $(1+x)_n = 1 + nx + n(n-1)x^2 + \dots$ $|x| < 1$ M1
Substitute in for $n = -3$
 $1 - 3x + \frac{(-3)(-4)x^2}{2} + \frac{(-3)(-4)(-5)x^3}{6}$ M1
 $= 1 - 3x + 6x^2 - 10x^3$ A1A1A1
(5)
- b) $(4 + 4x)^{-3} = 4^{-3} (1+x)^{-3}$ M1
 $\therefore (4 + 4x)^{-3} = \frac{1}{64} [1 - 3x + 6x^2 - 10x^3]$ A1ft (2)

4. a) $t = x - 1$ M1
Substitute in
 $\therefore y = (x-1)^2 + 2$ $[= x^2 - 2x + 3]$ M1A1(3)
- b) Using chain rule $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$, M1
 $\frac{dx}{dt} = 3\sin^2 t \cos t$ A1
Using $\frac{dt}{dx} = \frac{1}{\frac{dx}{dt}}$ M1
 $\therefore \frac{dy}{dx} = \frac{1 - \sin t}{3\sin^2 t \cos t}$ A1 (4)

c) i) Gradient = $\frac{dy}{dx}$, substitute in $t = \pi/4$
 $= (1 - \sin^2(\pi/4))/(3\sin^2(\pi/4)\cos^2(\pi/4))$ M1
 $= \frac{2\sqrt{2}-2}{3} = 0.27614\dots = 0.276$ (3 s.f.) A1 (2)

ii) gradient of normal at $t = \pi/4$ is $\frac{3}{2-2\sqrt{2}} = -3.6213\dots = -3.62$ (3 s.f.) M1 ft

$y = \frac{3}{2-2\sqrt{2}}x + c$ ($y = -3.62x + c$) M1

When $t = \pi/4$ ($=0.785\dots$), $x = 1 + \pi/4$ ($=1.785\dots$), $y = \pi/4 + \cos \pi/4$ ($= 1.492\dots$) A1
 Substitute in

$\pi/4 + \cos \pi/4 = \frac{3}{2-2\sqrt{2}}(1 + \pi/4) + c$

$c = \pi/4 + \cos \pi/4 - \frac{3}{2-2\sqrt{2}}(1 + \pi/4)$ ($= 7.9580\dots = 7.96$ (3 s.f.)) M1 ft

Therefore equation of normal is $y = \frac{3}{2-2\sqrt{2}}(x - 1 - \pi/4) + \pi/4 + \cos \pi/4$ A1 (5)

or $y = -3.62x + 7.96$ (3 s.f.)

5. a) $\frac{7}{(x+6)(2x+1)} = \frac{A}{x+6} + \frac{B}{2x+1}$ M1

$\therefore 7 = A(2x+1) + B(x+6)$ A1

Use $x = -6$

$\therefore 7 = -11A \therefore A = -\frac{7}{11}$ M1A1

Use $x = -\frac{1}{2}$

$\therefore 7 = B \times \frac{11}{2} \therefore B = \frac{14}{11}$ M1A1

$\therefore \frac{7}{(x+6)(2x+1)} = \frac{14}{11(2x+1)} - \frac{7}{11(x+6)}$ A1 (7)

b) $f'(x) = \frac{7}{11(x+6)^2} - \frac{28}{11(2x+1)^2}$ A1A1

$f'(1) = \frac{7}{539} - \frac{28}{99} = -\frac{17}{63}$ M1A1(4)

6. a) $r_1 - r_2$ M1

$\overline{AB} = 2\mathbf{i} + 0\mathbf{j} - 6\mathbf{k}$ A1A1(3)

b) Calculate Modulus = $\sqrt{2^2 + 6^2} = \sqrt{40}$ M1

$\frac{1}{\sqrt{40}}[2\mathbf{i} - 6\mathbf{k}]$ or $\frac{1}{\sqrt{10}}[\mathbf{i} - 3\mathbf{k}]$ A1 (2)

c) $-\mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda[2\mathbf{i} - 6\mathbf{k}]$ or $\mathbf{i} + 2\mathbf{j} - 5\mathbf{k} + \lambda[2\mathbf{i} - 6\mathbf{k}]$ A1A1(2)

d) Using $\mathbf{a} \cdot \mathbf{b} = ab\cos\theta = a_1b_1 + a_2b_2 + a_3b_3$ M1

Substitute in $\sqrt{40}\sqrt{30}\cos\theta = 2 + 30 = 32$ A1A1

$$\therefore \cos\theta = \frac{32}{20\sqrt{3}} \quad \text{M1}$$

$$\therefore \theta = 22.517\dots = 22.5^\circ \text{ (3 s.f.)} \quad \text{A1 (5)}$$

7. a) $\int \sin x \cos x \, dx = \frac{1}{2} \int \sin 2x \, dx \quad \text{M1}$

$$= \frac{1}{2} \times \frac{-\cos 2x}{2} = -\frac{\cos 2x}{4} \quad \text{A1}$$

Apply limits = $[0] - \left[-\frac{1}{4}\right] = \frac{1}{4} \quad \text{M1A1(4)}$

b) $I = \int e^x \sin x \, dx$, where $u = \sin x \quad v' = e^x$

$$u' = \cos x \quad v = e^x \quad \text{M1M1}$$

$$\therefore I = e^x \sin x - \int e^x \cos x \, dx \quad \text{A1}$$

Let $J = \int e^x \cos x \, dx$ and repeat integration, where $u = \cos x \quad v' = e^x$

$$u' = -\sin x \quad v = e^x \quad \text{M1}$$

$$J = e^x \cos x - \int -e^x \sin x \, dx$$

$$= e^x \cos x + I$$

$$\therefore I = e^x \sin x - [e^x \cos x + I] \quad \text{M1}$$

$$\therefore I = \frac{1}{2} e^x (\sin x - \cos x) + \mathbf{k} \quad \text{A2A1(8)}$$

c) $x \sin x = y \cos y$

$$\frac{d}{dx}(x \sin x) = \frac{dy}{dx} \times \frac{d}{dy}(y \cos y) \quad \text{M1}$$

attempt to use product rule M1

$$\sin x + x \cos x = \frac{dy}{dx} [\cos y - y \sin y] \quad \text{A1A1}$$

$$\therefore \frac{dy}{dx} = \frac{\sin x + x \cos x}{\cos y - y \sin y} \quad \text{A1 (5)}$$

Mark Scheme 3	Matching the syllabus written by EDEXCEL Curriculum 2004+
Calculators Allowed <i>Where appropriate leave your answers to 3 s.f.</i>	
© ZigZag Education 2004	Core Mathematics – C4

1. a) The curve is an *inverted exponential* which crosses the y-axis at $y = 1$ and the x-axis at $x = \ln(3/2) \approx 0.405\dots$ M1A1
A1 (3)
- b)
$$\int_2^3 (3 - 2e^x) dx$$

$$= [3x - 2e^x]_2^3$$
Substituting in limits;

$$= (9 - 2e^3) - (6 - 2e^2) = 3 - 2e^3 + 2e^2 = -22.392\dots$$

$$\therefore \text{Area} = -22.392\dots = 22.4 \text{ (3 s.f.) below x-axis}$$
 M1
A1A1
M1
A1 (5)
-
2. a) Use binomial theorem with suitable substitution M1
- $$(1 - 4x)^{1/2} = 1 + \left(\frac{1}{2}\right)(-4x) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2!}(-4x)^2 + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3!}(-4x)^3$$
- $$= 1 - 2x - 2x^2 - 4x^3$$
- A1
A1A1A1(5)
- b)
$$\left(1 - 4 \times \frac{1}{100}\right)^{1/2} = \left(100 - \frac{4}{100}\right)^{1/2} = \frac{\sqrt{96}}{10}$$

$$= \frac{4\sqrt{6}}{10}$$

$$\frac{4\sqrt{6}}{10} = 1 - 2 \times \frac{1}{100} - 2 \times \frac{1}{10000} - 4 \times \frac{1}{1000000} = 0.979796$$

$$\therefore \sqrt{6} = \frac{10}{4} \times 0.979796$$

$$\therefore \sqrt{6} = 2.44949 = 2.4495 \text{ (5 s.f.)}$$
 M1
A1
M1
A1 (4)
-
3. a) Using chain rule $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$, where $\frac{dx}{dt} = 2\sin t \cos t$ and $\frac{dy}{dt} = -\sin t$ M1A1A1
- $$\therefore \frac{dy}{dx} = \frac{-\sin t}{2\sin t \cos t} = -\frac{1}{2\cos t}$$
- A1 (4)
- b)
$$-\frac{1}{2\cos \pi/4} = A\sqrt{2}$$

$$-\frac{1}{2\sqrt{2}} = A\sqrt{2}$$

$$\therefore A = -\frac{1}{2}$$
 M1
A1 (2)
- c) $x = \sin^2 t$ $y = 1 + \cos t$
 $x = 1 - \cos^2 t$ $\therefore \cos t = y - 1$
 $\cos^2 t = (y - 1)^2$
Using $\sin^2 t + \cos^2 t = 1$
 $x = 1 - (y - 1)^2$ or further simplified i.e. $x = 2y - y^2$ M1
A1
A1 (3)
-
4. a) $u = \cos x$

$$\frac{du}{dx} = -\sin x \quad \text{A1}$$

$$\therefore dx = -\frac{1}{\sin x} du$$

$$\sin x \cos^3 x \, dx \Rightarrow \int u^3 \sin x \times \left(-\frac{1}{\sin x}\right) du \quad \text{M1}$$

$$= \int -u^3 \, du \quad \text{A1}$$

$$= -\frac{1}{4}u^4 + c \quad \text{A1}$$

$$= -\frac{1}{4}\cos^4 x + c \quad \text{A1 (5)}$$

$$\text{b) } \sin A + \sin B \equiv 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right) \quad \text{M1}$$

$$\frac{A+B}{2} = 6x \Rightarrow A+B = 12x$$

$$\frac{A-B}{2} = 5x \Rightarrow A-B = 10x$$

$$\therefore 2A = 22x \quad \text{M1}$$

$$A = 11x$$

$$B = x$$

$$\text{Thus } 2\sin x \cos 5x = \sin 11x + \sin x \quad \text{A1 (3)}$$

$$\text{c) } \int \sin 6x \cos 5x \, dx$$

$$= \int \frac{1}{2}(\sin 11x + \sin x) \, dx \quad \text{M1}$$

$$= -\frac{1}{22}\cos 11x - \frac{1}{2}\cos x + c \quad \text{A1A1(3)}$$

$$5. \text{ a) } \frac{x+1}{(x-1)^2(x+2)} \equiv \frac{A}{(x-1)} + \frac{B}{(x-1)^2} + \frac{C}{(x+2)} \quad \text{M1}$$

$$x+1 = A(x-1)(x+2) + B(x+2) + C(x-1)^2$$

$$\text{When } x = 1, 2 = B \times 3 \quad \text{M1}$$

$$\therefore B = \frac{2}{3} \quad \text{A1}$$

$$\text{When } x = -2, -1 = C \times (-3) \times 2$$

$$\therefore C = -\frac{1}{9} \quad \text{A1}$$

$$\text{Since } 0 = A + C$$

$$A = \frac{1}{9} \quad \text{A1}$$

$$\therefore \frac{x+1}{(x-1)^2(x+2)} = \frac{1}{9(x-1)} + \frac{2}{3(x-1)^2} - \frac{1}{9(x+2)} \quad \text{A1 (6)}$$

$$\text{b) } \frac{d}{dx} f(x) = -\frac{1}{9(x-1)^2} - \frac{4}{3(x-1)^3} + \frac{1}{9(x+2)^2} \quad \text{A1}$$

$$\frac{d}{dx} f(2) = -\frac{1}{9 \times 1} - \frac{4}{3 \times 1} + \frac{1}{9 \times 16}$$

$$= -1\frac{7}{16} \quad \text{A1 (2)}$$

c) Area = $\int_4^5 f(x) dx$ M1

$$\int_4^5 \left[\frac{1}{9(x-1)} + \frac{2}{3(x-1)^2} - \frac{1}{9(x+2)} \right] dx$$

$$= \frac{1}{9} \ln|x-1| - \frac{2}{3(x-1)} - \frac{1}{9} \ln|x+2| \Big|_4^5$$
A1A1A1

$$= \left(\frac{1}{9} \ln 4 - \frac{1}{6} - \frac{1}{9} \ln 7 \right) - \left(\frac{1}{9} \ln 3 - \frac{2}{9} - \frac{1}{9} \ln 6 \right)$$

$$= \frac{1}{9} \ln \left(\frac{4}{7} \right) - \frac{1}{6} - \frac{1}{9} \ln \left(\frac{1}{2} \right) + \frac{2}{9}$$


$$= \frac{1}{9} \ln \left(\frac{8}{7} \right) + \frac{1}{18}$$
A1 (5)

6. a) $\int_1^4 e^x dx = [e^x]_1^4 = e^4 - e^1 = 51.8799.. = 51.9$ or better **correctly substituting limits** M1A1A1(3)

b) i) $h = 1$; $y_0 = e^{-1}$; $y_1 = e^{-2}$; $y_2 = e^{-3}$; $y_3 = e^{-4}$; M1M1

Area = $\frac{1}{2} [e^{-1} + e^{-4} + 2(e^{-2} + e^{-3})] = 0.37821... = 0.378$ (3 s.f.) M1A1(4)

ii) In each interval an estimate of the area of the region is obtained by joining successive points on the curve with a straight line. Any two points connected form a region which includes **a section above the line**. Thus the area calculated by the trapezium rule is an **over estimate**. Suitable diagram.



M1

To improve accuracy, need to take more intervals. A1 (2)

c) These two graphs are reflections of each other in the y-axis A1 (1)

7. a) $\vec{AB} = (i + 2j + k) - (-i - 2j + k)$ M1

$\vec{AB} = 2i + 4j$ A1 (2)

b) Magnitude = $\sqrt{(-1)^2 + (-2)^2 + 1^2}$ M1A1

$$= \sqrt{1+4+1}$$

$$= \sqrt{6}$$
A1 (3)

$\vec{OA} = \frac{1}{\sqrt{6}} (-i - 2j + k)$

c) Distance = $\sqrt{2^2 + 4^2}$ M1

$$= \sqrt{20} \text{ units}$$

$$= 2\sqrt{5} \text{ units}$$
A1 (2)

d) $i + \mu 3i = 3i + \lambda i$

$$\mathbf{i}(1 + 3\mu) = \mathbf{i}(3 + \lambda)$$

$$\therefore 1 + 3\mu = 3 + \lambda \quad \leftarrow$$

M1

$$2\mathbf{j} + \mu 2\mathbf{j} = 2\mathbf{j} + \lambda 2\mathbf{j}$$

$$\mathbf{j}(2 + 2\mu) = \mathbf{j}(2 + 2\lambda)$$

M1

$$\therefore \mu = \lambda$$

Substitute into \leftarrow

$$1 + 3\lambda = 3 + \lambda$$

$$\therefore 2\lambda = 2$$

$$\lambda = 1 \Rightarrow \mu = 1$$

A1

$$\therefore (3 + \mu)\mathbf{k} = (1 + A\lambda)\mathbf{k}$$

$$\therefore 3 + 1 = 1 + A$$

$$\therefore A = 3$$

A1 (4)

e) $a_1b_1 + a_2b_2 + a_3b_3 = (1)(3) + (2)(2) + (1)(3)$
 $= 10$

M1

A1

$$|a| = \sqrt{9+4+1} = \sqrt{14}$$

$$|b| = \sqrt{1+4+9} = \sqrt{14}$$

$$14\cos\theta = 10$$

B1

$$\cos\theta = \frac{5}{7} = 44.415\dots = 44.4^\circ \text{ (3 s.f.)}$$

A1 (4)

(75)


Mark Scheme 4	Matching the syllabus written by EDEXCEL Curriculum 2004+
Calculators Allowed <i>Where appropriate leave your answers to 3 s.f.</i>	
© ZigZag Education 2004	Core Mathematics – C4

1. $x^2 = 9\cos^2 2t$ $y^2 = 9\sin^2 2t$
squaring and adding M1
 $\therefore x^2 + y^2 = 9\cos^2 2t + 9\sin^2 2t = 9(\cos^2 2t + \sin^2 2t) = 9$ M1A1
 \therefore **Circle**, with centre (0, 0), radius 3 A1A1A1(6)
-
2. $\int dy = \int (x + \sin x) dx$ M1
 $y = \int (x + \sin x) dx$ A1
 $= \frac{x^2}{2} - \cos x + c$ A1
Sub in 0.1 and 0.2 to work out c M1
 $\therefore y = \frac{x^2}{2} - \cos x + 1.1900\dots = 1.19$ (3 s.f.) A1 (5)
-
3. a) Using binomial expansion M1
 $(1 - 2x)^{\frac{1}{2}} = 1 + \left(\frac{1}{2}\right)(-2x) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2!}(-2x)^2 + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3!}(-2x)^3$ M1
 $= 1 - x - \frac{1}{2}x^2 - \frac{1}{2}x^3$ A1A1(4)
- b) Sub in: $1 - 0.01 - \frac{1}{2}(0.01)^2 - \frac{1}{2}(0.01)^3 = 0.9899495$ M1
equating $\therefore \left(1 - \frac{2}{100}\right)^{\frac{1}{2}} = 0.9899495$ M1
 $\therefore \frac{1}{10}(98)^{\frac{1}{2}} = 0.9899495$
 $\therefore \frac{7}{10}\sqrt{2} = 0.9899495$ M1
 $\therefore \sqrt{2} = \left(\frac{10}{7}\right) \times 0.9899495 = 1.41421\dots = 1.4142$ (5 s.f.) A1 (4)
-
4. a) Integration by parts M1
 $I = \int x \sin x dx$ $u = x$ $v' = \sin x$ M1A1
 $u' = 1$ $v = -\cos x$
 $\therefore I = -x \cos x - \int (-\cos x) \times 1 dx$
 $= -x \cos x + \sin x + c$ A1A1(5)
- b) $I_2 = \int x^2 \cos x dx$ $u = x^2$ $v' = \cos x$ M1A1
 $u' = 2x$ $v = \sin x$
 $I_2 = x^2 \sin x - \int 2x \sin x dx = x^2 \sin x - 2I$ M1

$$= x^2 \sin x + 2x \cos x - 2 \sin x + k$$

A1A1(5)

5. a) $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$ M1
 $\frac{dy}{dt} = -2 \cos t \sin t, \quad \frac{dx}{dt} = -6 \sin 6t$ A1A1
 Using $\frac{dt}{dx} = \frac{1}{\frac{dx}{dt}}$ M1
 $\frac{dy}{dx} = \frac{-2 \cos t \sin t}{-6 \sin 6t} = \frac{\cos t \sin t}{3 \sin 6t}$ A1 (5)
- b) Substitute into equation above to find the gradient M1
 i.e. Gradient = $\frac{\frac{1}{\sqrt{2}} \times \frac{1}{\sqrt{2}}}{3 \times -1} = -\frac{1}{6}$ A1 (2)
- c) $m_1 m_2 = -1 \quad \therefore m_2 = 6$ A1 ft
 $y = 6x + c$
 Sub in t to find that $x = 0$ and $y = \frac{1}{2}$ and sub into $y = 6x + c$ M1
 $\therefore c = \frac{1}{2} \quad \therefore y = 6x + \frac{1}{2}$ A1 (3)

6. a) $\int \frac{x^3}{x^4+1} dx = \ln|x^4+1| + k$ OR $\ln|k(x^4+1)|$ (A1 for missing k) A2 (2)
- b) $\int_1^5 x^{1/2} + e^x dx = \left[\frac{2x^{3/2}}{3} + e^x \right]_1^5$ A1A1
 Substitute in limits correctly; $= \left[\frac{2 \times 5^{3/2}}{3} + e^5 \right] - \left[\frac{2}{3} + e \right]$ M1
 $= 155.86... - 3.3849...$
 $= 152.48... = 152.5$ (1 d.p.) A1 (4)
- c) $h = 1, n = 4 \Rightarrow A = \frac{1}{2}(y_0 + 2(y_1 + y_2 + y_3) + y_4)$ M1
 $\int_1^5 \frac{1}{x^{1/2} + e^x} dx \approx \frac{1}{2}(f(1) + 2(f(2) + f(3) + f(4)) + f(5))$ M1
 $= \frac{1}{2}(0.26894... + 2(0.11359... + 0.045834... + 0.017668...) + 0.0066379...)$ A1
 $= 0.31488... = 0.315$ (3 s.f.) A1
 The trapezium rule gives an **over**-estimate of the integral. A1
 Shading between curve and line segment.  M1 ft(6)

7. a) $a \cdot b = |a||b| \cos \theta = a_1 b_1 + a_2 b_2 + a_3 b_3$ M1
 $|a| = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3} \quad |b| = \sqrt{3^2 + 2^2 + 2^2} = \sqrt{17}$ A1

- $\therefore \cos \theta = \frac{-3+2+2}{\sqrt{3}\sqrt{17}} = 0.14002\dots$ A1
 $\therefore \theta = 81.9505\dots^\circ = 81.951$ (3 d.p.) A1 (4)
- b) $|AB| = \sqrt{4^2 + 1^2 + 1^2} = \sqrt{18}$ M1A1(2)
- c) $\vec{AC} = (c+1)i + 4k$ $\vec{AB} = 4i + j + k$ A1A1
 $(\vec{AC}) \cdot (\vec{AB}) = 4(c+1) + 0 + 4 = 0$ M1A1 ft
 $4c + 8 = 0 \quad \therefore c = -2$ A1 (5)
- d) $-i + j + k$ (or other valid position) $+ \lambda(4i + j + k)$ (or multiple, including -1) A1A1(2)
-

8. a) $f(x) = \frac{12x}{(x+1)(2x+1)} = \frac{A}{x+1} + \frac{B}{2x+1}$ M1
 $12x = A(2x+1) + B(x+1)$
 $x = -1 \Rightarrow -12 = -A \Rightarrow A = 12$ $x = -0.5 \Rightarrow -6 = 0.5B \Rightarrow B = -12$
 $\therefore f(x) = \frac{12}{x+1} - \frac{12}{2x+1}$ A1A1(3)
- b) $f'(x) = -\frac{12}{(x+1)^2} + \frac{24}{(2x+1)^2}$ A1ftA1ft
(2)
- c) $f(x) = 12[(x+1)^{-1} - (2x+1)^{-1}]$ M1
 $(x+1)^{-1} = 1 - x + x^2 + \dots \quad -1 < x < 1$ A1
 $(2x+1)^{-1} = 1 + (-1)(2x) + \frac{(-1)(-2)}{2}(2x)^2 + \dots$ M1
 $= 1 - 2x + 4x^2 + \dots \quad -0.5 < x < 0.5$ A1
 $\therefore f(x) = 12[x - 3x^2]$ A1A1(6)
-

Mark Scheme 5	Matching the syllabus written by EDEXCEL Curriculum 2004+
Calculators Allowed <i>Where appropriate leave your answers to 3 s.f.</i>	
© ZigZag Education 2004	Core Mathematics – C4

1. Using $V = \pi \int y^2 dx$ M1
- $$y^2 = \left(e^{\frac{x}{2}}\right)^2 = e^x$$
- A1
- $$= \pi \int_2^3 e^x dx$$
- $$= \pi [e^x]_2^3$$
- A1
- Substitute in limits M1
- $$= \pi (e^3 - e^2)$$
- A1
- (5)
-
2. Separate variables $\Rightarrow \int dy = \int (x + e^x) dx$ M1
- $$\therefore y = \int (x + e^x) dx \quad \Rightarrow \quad y = \frac{x^2}{2} + e^x + c$$
- A1A1
- Sub in $x = 1, y = 2$ M1
- $$\therefore c = \frac{3}{2} - e = -1.2182... = -1.22 \text{ (3 s.f.)}$$
- A1 (5)
-
3. a) $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$ M1
- $$\frac{dy}{dt} = 3 \cos 3t, \quad \frac{dx}{dt} = \cos t$$
- A1A1
- $$\therefore \frac{dy}{dx} = \frac{3 \cos 3t}{\cos t}$$
- A1 (4)
- b) Sub in $t = \frac{\pi}{4}$ M1
- $$\therefore \frac{dy}{dx} = -3$$
- A1 (2)
- c) $y = -3x + c$ M1
- When $t = \frac{\pi}{4}, x = \frac{1}{\sqrt{2}}, y = \frac{1}{\sqrt{2}}$ M1
- $$\therefore \frac{1}{\sqrt{2}} = \frac{-3}{\sqrt{2}} + c \quad \therefore c = \frac{4}{\sqrt{2}} \text{ or } 2\sqrt{2} (= 2.8284... = 2.83 \text{ (3 s.f.)})$$
- A1 (3)
-
4. a) $I = \int (3 \sin 6x - \sin 2x) dx = -\frac{1}{2} \cos 6x + \frac{1}{2} \cos 2x + C$ M1A1A1A1
- (4)
- b) $u = \sin x \quad \frac{du}{\cos x} = dx$ sub this in M1 ft
- $$\therefore I = 16 \int u^4 du = \frac{16u^5}{5} = \frac{16 \sin^5 x}{5}$$
- A1A1
- Sub in limits $\therefore I = \frac{1}{10}$ A1 (4)
-

5. a) $D^2 = (b-1)^2 + (b+1)^2 + 12^2$ A1
 Equate to 14^2 M1
 $2b^2 + 146 = 196 \quad \therefore b = +5 \text{ or } -5$ A1A1(4)
- b) a. $c = 2 - 1 - 2 = -1$ A1
 $|a| = \sqrt{3}, |c| = \sqrt{9} = 3$ A1A1
 a. $c = (\mathbf{a})(\mathbf{c})\cos \theta = \underline{a_1c_1} + \underline{a_2c_2} + \underline{a_3c_3}$ M1M1
 $\cos \theta = \frac{-1}{3\sqrt{3}}$ A1
 \therefore Acute angle $= 180^\circ - \theta = 78.904\dots = 78.9^\circ$ (3 s.f.) A1 (7)
- c) $\mathbf{i + j + k}$ (or other correct position) $+ \lambda(\overrightarrow{AC})$ (or multiple including -1) A1A1
 $= \mathbf{i + j + k} + \lambda(\mathbf{i - 2j - 3k})$
 $\mathbf{i} = 17 \quad \therefore \lambda = 16$ M1
 for the \mathbf{j} we have: $1 + (-2 \times 16) = -31$ so consistent and so can pass through that point B1
 \therefore position is $17\mathbf{i} - 31\mathbf{j} - 47\mathbf{k}$ i.e. $D = 47$ A1 (5)
-
6. a) i) Multiply equations for same coefficients:
 $3x = 6\cos t$
 $2y = 6\sin t$ M1
 Square both sides to give
 $9x^2 = 36\cos^2 t$
 $4y^2 = 36\sin^2 t$ M1
 $9x^2 + 4y^2 = 36\cos^2 t + 36\sin^2 t$ M1
 Using $\sin^2 t + \cos^2 t = 1$, Cartesian equation is $9x^2 + 4y^2 = 36$
 M1A1(5)
- ii) Differentiate implicitly M1
 $18x + 8y \frac{dy}{dx} = 0$ A1A1A1
 $\frac{dy}{dx} = -\frac{9x}{4y}$ A1 (5)
- b) i) $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx} \quad \left[\frac{dt}{dx} = \frac{1}{\left(\frac{dx}{dt}\right)} \right]$ M1M1
 $\frac{dy}{dt} = 3\cos t, \frac{dx}{dt} = -2\sin t$
 $\frac{dy}{dx} = -\frac{3\cos t}{2\sin t}$ or $-\frac{3}{2}\cot t$ A1 (3)
- ii) $\frac{dy}{dx} = -\frac{3 \times \frac{x}{2}}{2 \times \frac{y}{3}} = -\frac{9x}{4y}$, same as aii). M1A1(2)
-
7. a) $\frac{36x}{(2x+1)(x+2)} \equiv \frac{A}{2x+1} + \frac{B}{x+2}$ M1
 $36x \equiv A(x+2) + B(2x+1)$ M1
 Let $x = -2 \quad \therefore -72 = -3B \quad B = 24$ A1
 Let $x = 0 \quad \therefore 0 = 2A + B \quad A = -12$ M1A1
 $\therefore \frac{36x}{(2x+1)(x+2)} = -\frac{12}{2x+1} + \frac{24}{x+2}$ A1 (6)

b) $f(x) = 12[-(2x + 1)^{-1} + 2(x + 2)^{-1}]$ M1
Using binomial expansion, M1

$$(2x + 1)^{-1} = 1 + (-1)(2x) + \frac{(-1)(-2)(2x)^2}{2} + \dots$$

M1

$$= 1 - 2x + 4x^2 + \dots \quad -1 < 2x < 1$$
A1

$$(x + 2)^{-1} = (2^{-1})\left(\frac{x}{2} + 1\right)^{-1}$$
M1

$$\left(\frac{x}{2} + 1\right)^{-1} = 1 + (-1)\left(\frac{x}{2}\right) + \frac{(-1)(-2)\left(\frac{x}{2}\right)^2}{2} + \dots = 1 - \frac{x}{2} + \frac{x^2}{4} \quad -1 < \frac{x}{2} < 1$$
M1A1B1

$$\therefore f(x) = 12[-(1 - 2x + 4x^2) + \frac{1}{2} \times 2\left(1 - \frac{x}{2} + \frac{x^2}{4}\right)] = 12\left(\frac{3}{2}x - \frac{15}{4}x^2\right) = \mathbf{18x - 45x^2}$$
A1A1

for $-\frac{1}{2} < x < \frac{1}{2}$ A1 (11)

(75)