

## Mark Scheme (Results)

January 2013

International GCSE Chemistry (4CH0) Paper 1C Science Double Award (4SC0) Paper 1C

Edexcel Level 1/Level 2 Certificate Chemistry (KCH0) Paper 1C Science (Double Award) (KSC0) Paper 1C





## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at <u>www.edexcel.com</u> or <u>www.btec.co.uk</u> for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at <u>www.edexcel.com/contactus</u>.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson. Their contact details can be found on this link: <u>www.edexcel.com/teachingservices</u>.

You can also use our online Ask the Expert service at <u>www.edexcel.com/ask</u>. You will need an Edexcel username and password to access this service.

## Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: <a href="https://www.pearson.com/uk">www.pearson.com/uk</a>

January 2013 Publications Code UG034348 All the material in this publication is copyright © Pearson Education Ltd 2013

	iesti			Answer	Notes	Marks
ทเ	number					
1	а		M1	nucleus		1
			M2	protons	Accept in either order	1
			M3	neutrons	Accept in either order	1
			M4	electrons		1
			M5	shells		1
			M6	protons AND electrons	In either order	1
			M7	electrons		1
	b	i		3		1
		ii		5		1
2	а	i		С		1
		ii		В		1
	b			fluorine / F <sub>2</sub>	Accept F	1
	С	i		hydrogen chloride		1
		ii		hydrochloric (acid)		1
		iii		HCI		1
					Total	15

	uest umb			Answer	Notes	Marks
3	а	i	M1	bubbles / fizzing / effervescence	Accept gas formed /given off	1
					Ignore any name or formula	
			M2	iron/solid disappears	Accept iron/solid gets smaller /	1
				OR	dissolves	
				green/colourless solution (forms)		
					Ignore references to heat change / change in pH	
		ii		iron sulfate AND hydrogen (in either order)	Penalise oxidation states other than (II)	1
					Accept ferrous sulfate	
					Reject ferric sulfate	
	b			2 (1) 2	Accept multiples and fractions	1
	С	i	M1	white	Ignore colourless	1
			M2	blue	Ignore all qualifiers such as pale / dark	1
		ii		D		1
		iii	M1	boiling point / melting/freezing point		1
			M2	100 °C / 0 °C	Accept just ° or C but not just number	1
					Value must match property	
					Accept correct values in K	
					Ignore other physical properties such as pH / density	

	Question number		Answer	Notes	Marks
3	d	i	low density / less dense than air	Accept lighter than air / the lightest gas but not just light / lightweight	1
		ii	non-flammable OR does not burn / explode (when ignited)	Ignore unreactive Accept does not react with oxygen/air	1
	e		$H_2(g) \rightarrow H_2(I)$	Ignore "+ heat/energy" on RHS Penalise indication of endothermic process	1
				Total	12

	Question number			Answer	Notes	Marks
4	а	i		oxygen / air	Accept O <sub>2</sub> but not just O	1
		ii		iron(III) oxide / ferric oxide	Ignore hydrated	1
					Accept iron oxide but not ferrous oxide	
					or iron oxide with an incorrect	
					oxidation state	
	b	i		nail is wet / needs drying		1
				OR		
				rust absorbs water/is wet / rust falls off	Not just some of the nail/it falls off	
		ii		bigger/biggest increase/change in mass	Accept mass increased more	1
				OR	Ignore highest mass	
				mass of rust greater/greatest	Ignore just mass increased by 0.3 g	
					Accept weight in place of mass	
		iii		bigger/biggest increase/change in mass with		1
				correct reference to proportion or percentage		
		iv		mass has decreased	Accept mass has not increased /	1
					should have increased	
					Accept mass cannot decrease	
					Accept nail got lighter	
	С		M1	grease / oil		1
			M2	paint		1

	Question number			Answer	Notes	Marks
4	d		M1	galvanising / sacrificial (protection)	Ignore references to anode / cathode	1
			M2	zinc more reactive (than iron) OR iron less reactive (than zinc)	Accept zinc higher in (re)activity series OR iron lower in (re)activity series Must be comparison, eg not just zinc is reactive without statement that iron is unreactive	1
			M3	zinc reacts/corrodes/oxidises instead of iron OR zinc reacts/corrodes/oxidises before iron	Accept zinc reacts faster than iron Accept zinc loses electrons instead of/before iron / zinc gives electron(s) to iron (ions) Reject zinc rusts Reject protective layer of zinc oxide Ignore erodes	1
					Total	11

	Question number			Answer	Notes	Marks
5	а		M1	(compound/molecule/substance containing) carbon and hydrogen (atoms)	Reject atoms/elements in place of compounds Reject molecules in place of atoms Reject mixture Accept C and H in place of carbon and hydrogen	1
			M2	only	M2 dependent on M1 or near miss, eg mixture of C and H Accept equivalent wording such as alone / purely / solely	1
	b			contains (C=C) double bonds	Accept multiple bonds Reject implied C=H	1

	Question number		Answer		Notes	Marks
5	С	i		alkene(s)		1
		ii		C <sub>n</sub> H <sub>2n</sub>	Accept other symbols such as x Accept H <sub>2n</sub> C <sub>n</sub>	1
		111	M1	same/similar chemical properties	Accept same/similar reactions Do not accept a specific reaction, eg they all burn Ignore similar reactivities	
			M2	trend/gradation in physical properties	Accept named trend eg boiling point Accept correct trend eg smaller molecules have lower boiling points, but not incorrect trend such as smaller molecules have higher boiling points	2
			M3	same functional group		
	-		M4	(neighbouring) members differ by CH <sub>2</sub>		
l					Any two for 1 each	

	Question number												Answer	Notes	Marks
5	d	i			Accept butene Ignore mention of cis or trans	1									
		ii		C <sub>4</sub> H <sub>8</sub>		1									
		iii	M1	(compounds/molecules with) same molecular formula / same number of each type of atom	Do not penalise specific compound types, eg hydrocarbons / alkenes If elements/atoms in place of compounds, max 1 for Q Ignore references to chemical/general/empirical formula	1									
			M2	different structure(s) / different structural formula(e) / different displayed formula(e)	Ignore atoms in a different order	1									
		iv		displayed formula of but-2-ene or methylpropene	Accept cyclobutane or methylcyclopropane Ignore but-1-ene structure	1									
	е	i		colourless / decolorised	Ignore clear	1									
		ii		C <sub>2</sub> H <sub>4</sub> Br <sub>2</sub>	Insist on correct use of subscripts and cases of letters Do not penalise elements in different order Accept correct structural/displayed formula	1									
					Total	14									

	estion Imber		Answer	Notes	Marks
6	а	M1 M2	(same) volume of solution/liquid (same) concentration (of solution/liquid)	Accept amount in place of volume	
		M3	(same) amount of metal	Accept solid in place of metal Accept moles/mass in place of amount Reject salt in place of metal	3
		M4	(same) (total) surface area / state of subdivision of solid	Accept all powder / all particles same size / size of metal	
		M5	(same) method/length of time/speed of stirring		
				Ignore references to polystyrene cup / starting temperature	
				Any three for 1 each	
	b	M1	22.4	Ignore trailing zeroes and award 1 mark	1
		M2	17.7	for both correct values in wrong order	1
		M3	4.7	Consequential on values used in M1 and M2 Reject –4.7	1

	Question number			Answer	Notes	Marks
6	С	i	M1	magnesium / Mg		1
			M2	larg <u>est</u> difference in (recorded) temperatures	DEP on M1 Do not accept results in place of temperatures Explanation must be comparative: Not just rises are 10.5 and 15.5 without reference to values for other metals Not just very different temperature changes Accept two temperatures furthest apart Not just largest temperature rise No penalty for quoting wrong difference, eg 4 C	1

	Question number			Answer	Notes	Marks
6	С	ii	M1	magnesium / Mg		1
			M2	larg <u>est</u> temperature <u>rise/change</u>	DEP on M1 Do not accept results in place of temperatures Accept answers stating most heat produced / most exothermic	1
		III		silver/it is less reactive (than copper) / lower in the reactivity series OR no reaction	Must be implied comparison Accept copper more reactive than silver but not just silver is unreactive	1
					Reject references to differences in reactivities of silver and copper ions / silver and copper sulfate	
		iv		silver and X both have no temperature rise/change OR two metals show no temperature rise/change	Accept two metals did not react Accept two showed zero (temperature rises)	1
	d			$Zn + CuSO_4 \rightarrow Cu + ZnSO_4$	Ignore state symbols Accept correct ionic equation with or without spectator ions	1
					Total	13

	iesti			Answer	Notes	Marks
n	<u>umb</u>	er				
7	а	i	M1	(A) reduced AND (B) oxidised	If first column blank, M1 can be scored from words in second column: eg reduction is gain of electrons in 2nd column for A scores M2 oxidation is gain of oxygen in 2nd column for B scores M3 Both above statements would score M1 as well If M1 wrong way round, then 0/3 If 1st column has oxidation ×2 or reduction × 2, then no M1, but it will sometimes be possible to award either M2 or M3 for a correct	1
					matching statement.	
			M2	(A) gain of electrons	If species identified, must be correct,	1
					eg not aluminium gains electrons	
					Ignore reference to loss of oxygen	
			М3	(B) gain of oxygen	Ignore reference to loss of electrons	1
		ii	M1		Both formulae correct and on correct	1
				$2O^{2-} \rightarrow O_2 + 4e^{(-)}$	sides of equation	
			M2		Balancing including electrons	1
					Accept electrons lost on LHS	
					Accept $O^{2-} \rightarrow O + 2e^{(-)}$ followed by	
					$20 \rightarrow 0_2$	
		iii		electrodes burn /get smaller / decrease (in mass)	Ignore erode / corrode / wear /	1
				/ have to be replaced	damage	

	iest			Answer	Notes	Marks
ทเ	number					
7	а	iv	M1	coke	Ignore coal / carbon	1
					Reject other raw materials such as	
					limestone/haematite	
			M2	produces heat / exothermic (reaction)	M2 independent	1
					Accept makes carbon dioxide which	
					then produces CO/reducing agent	
	b	i		carbonating drinks / in drinks		1
			M2	soluble (in water) / reacts with water	M2 dependent on M1	1
					Accept just solubility, ignoring	
					qualifying statements such as only	
					slightly soluble, or only dissolves	
					under pressure	
			M3	fire extinguishers / putting out fires		1
			M4	denser than air / does not support combustion	Accept denser than oxygen	1
					Ignore does not burn	
					M4 dependent on M3	
					Ignore references to covering /	
					sitting on fire	
					Accept M1+M2, and M3+M4 reversed	

	Question number			Answer	Notes	Marks
7	b	ii	M1	$SO_2 + H_2O \rightarrow H_2SO_3$	Accept SO <sub>2</sub> + H <sub>2</sub> O + $\frac{1}{2}O_2 \rightarrow H_2SO_4$	1
					Do not accept unconventional	
					formulae such as SO <sub>3</sub> H <sub>2</sub>	
			M2	adverse effect on plants/trees/crops/vegetation OR	eg kills / harms / damages / destroys / stunts growth	1
				adverse effect on fish / water animals / aquatic life OR	eg kills / harms / damages / destroys /reduces fish population	
				adverse effect on iron/steel/metal OR	eg damages / causes rusting/corrosion	
				adverse effect on stonework/limestone/marble	eg causes corrosion / damages Not just buildings/structures/statues	
					Ignore changes in pH	
					Ignore effects on animals/birds Ignore just habitats	
					Do not accept	
					burning/weathering/erosion as	
					adverse effects	
					Not just affects plants/fish/etc	
					Do not apply list principle	

Question number		Answer	Notes	Marks
7 C	M1	$M_{\rm r}({\rm PbCO_3}) = 267 / A_{\rm r} ({\rm Pb}) = 207$	Accept 414 for Pb	1
	M2	$n(PbCO_3) / n(Pb) = 1.87 mol$	Accept 1.9	1
	M3	m(Pb) = 387 – 388 g		1
		OR		
	M1	$M_{\rm r}({\rm PbCO}_3) = 267 / A_{\rm r} ({\rm Pb}) = 207$		1
	M2	$m(Pb) = 207 \times 500$		1
		267		
	M3	m(Pb) = 387 - 388 g		1
		OR		
	M1	$M_{\rm r}({\rm PbCO}_3) = 267 / A_{\rm r} ({\rm Pb}) = 207$		1
	M2	% Pb = <u>207 × 100</u> = 77.5 %		1
		267		
	M3	m(Pb) = 500 × 0.775 = 387.5 g		1
			Correct final answer with or without	
			working scores 3	
			Accept answer in kg	
			Apply ECF throughout	
			Accept other valid methods	
			774-776 scores 2	
			Total	17

Question number			Answer		Notes	Marks	
8 8	a	i	M1	reversible (reaction) / goes forwards and/or backwa / can go in either direction	ards	Ignore equilibrium	1
			M2	enthalpy/heat/energy change		Ignore kJ/mol Reject energy produced/released	1
		ii		exothermic / heat/energy give	n out/lost	Accept enthalpy in place of heat/energy Ignore references to temperature	1
k	b		M1	two (vaguely) horizontal lines: one with reactants or their for AND one with products or their forr	mulae	Ignore all curves and connecting lines Ignore line representing <i>x</i> -axis and any label Accept R for reactants and P for products	1
			M2	reactants (line) above product	s (line)	<ul> <li>No penalty for products to left of reactants</li> <li>Accept formulae in place of words for reactants and products</li> <li>Do not penalise minor errors in</li> </ul>	1
						formulae (e.g. NH instead of NH <sub>3</sub> ) or missing coefficients	
(	С		M1	(effect of temp on rate)	increased		1
			M2	(effect of temp on yield)	decreased		1
			M3	(effect of catalyst on rate)	increased		1
			M4	(effect of catalyst on yield)	unchanged		1

	Question number			Answer	Notes	Marks
8	d	i	M1	decreased	No ECF from increased / no effect Accept longer time for reaction Ignore references to equilibrium	1
			M2	particles further apart/more widely spaced / more space to move in / concentration decreases	Accept molecules Reject atoms/ions in M2 only If neither of M2 and M3 scored, accept fewer collisions with no reference to frequency or time	1
			M3	less frequent (successful) collisions / fewer (successful) collisions per second/minute	Accept more time between collisions Ignore decreased chance / probability / likelihood of collisions References to change in energy/speed of particles means M2 and M3 cannot be scored	1
		ii	M1	shifted to right / more products / shifts in exothermic/forward direction	Ignore references to rate No ECF from shift to left / no change Accept forward reaction favoured	1
			M2	more (gas) moles/molecules on right	Accept fewer (gas) moles on left Accept favours side with more (gas) moles Accept 9 moles on left and 10 moles on right	1
	е			4 (1) 2 4	Accept fractions and multiples	1
					Total	15

	Question number			Answer	Notes	Marks
9	а	i	M1	35 on lines 1 and 3		1
			M2	44 on line 2		1
		ii		isotopes		1
		iii		same number of electrons (in outer shell) OR same electron arrangement or configuration	Ignore references to protons and neutrons unless incorrect, eg different numbers of protons, same number of neutrons	1
		iv	M1	<sup>79</sup> Br	Accept just 79	1
			M2	79 is closer to 79.9/more accurate value	Accept 79 is closer to relative atomic mass M2 dependent on M1	1

	Question number			Answer	Notes	Marks
9	b	i	M1		shared pair of electrons	1
			M2	××	other electrons correct (not necessary to be	1
				H*Br ×	paired)	
				Xx	M2 dependent on M1	
					Accept any combinations of dots and	
					crosses	
					Circles not needed but if drawn must	
					overlap or touch $-$ if not, then $0/2$	
					Ignore inner electron shells even if	
					incomplete or incorrect	
					Do not penalise incorrect symbols, eg br/BR	
					If Na used in place of H, max 1	
					No marks if ions shown	
		ii	M1	shared (two/pair of) electrons	Not share an electron	1
			M2	attracted to both nuclei	M2 dependent on M1 or near miss	1
					eg the electrons are attracted to the	
					nucleus scores 0	
					the electrons are attracted to both nuclei	
					scores M2 but not M1	
					0/2 if references to ions / ionic bond /	
					intermolecular forces	

	Question number			Answer	Notes	Marks
9	b	iii	M1	(sodium bromide) ionic bonding / + and – ions	Reject covalent bonding / shared electrons	1
			M2	<ul><li>(hydrogen bromide) attraction between</li><li>molecules</li><li>/ intermolecular forces (of attraction)</li></ul>	Accept dipole-dipole attractions / van der Waals' forces / IMF / vdW Ignore hydrogen bonds Reject ions/ionic	1
			M3	ionic bonding stronger OR IMF / attractions between HBr molecules weaker	Accept ionic bonds stronger M3 dependent on comparison of intermolecular forces and ionic bonding Accept correct references to energy needed to overcome bonding / attractions	1
					Ignore references to reactivity and mass	

	Question number				ŀ	Answer	Notes	Marks
9	C		M1	Na <u>13.8</u> 23	Br <u>47.9</u> 80	0 <u>38.3</u> 16	0/3 if division by atomic number(s) /division wrong way round If only two elements shown correctly, only M1 can be awarded	1
			M2 M3		0.6	2.4	Accept 1 : 1: 4Accept elements in any orderPenalise M3 for incorrect symbol, egSBrO4 or NaBO4Dividing by 160 instead of 80 givesNa2BrO8Dividing by 32 instead of 16 givesNaBrO2Award 2 in these casesBoth these errors give Na2BrO4Award 1 in this caseCorrect final answer scores 3 marks	1
							Total	16

Question number			Answer	Notes	Marks	
10	а			organic compounds flammable OR decreases chance of fire OR less vapour/gas escapes	Ignore references to breaking boiling tube / beaker/escape of mercury / need to hold boiling tube / being burned by flame / loss of heat Ignore liquid escapes Accept stops/prevents vapour escaping Reject references to reactions inside the beaker	1
10	b	i	M1 M2	all five points correct	to nearest gridline Deduct 1 mark for each error If points not visible, assume they are under the line	2
			M3	straight line of best fit	Must be drawn with a ruler Does not need to be extrapolated Line should go through any two correctly plotted points	1
		ii		correct qualitative relationship	eg boiling point increases as relative formula mass increases / positive correlation Accept statement "wrong" way round Reject mass in place of relative formula mass Reject temperature in place of boiling point Reject (directly) proportional	1
		iii		117 (°C) ±1°	CQ on candidate graph	1
		iv		E		1
					Total	7

Further copies of this publication are available from Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467 Fax 01623 450481 Email <u>publication.orders@edexcel.com</u>

Order Code UG034348 January 2013

For more information on Edexcel qualifications, please visit our website <u>www.edexcel.com</u>

Pearson Education Limited. Registered company number 872828 with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE





