
K.C.S.E CHEMISTRY PAPER 233/1 2001

Study the nuclear reactions given in the scheme below and answer the questions that follow.



- Student Bounty.com are isotopes. What is meant by the term isotope?
- (1 mark) Write an equation for the nuclear reaction in Step II
- Give one use of 14 C (1 mark)
- In an experiment, 0.8g of magnesium powder were reacted with excess dilute sulphuric acid at 25°C. The time for the reaction to come to completion was recorded. The experiment was repeated at 40°C. In which experiment was the time taken shorter? Explain your answer. (3 marks)
- The electronic structures for elements represented by letters A, B, C and D are D 2.8.8.
 - B 2.8.2; C 2.8.1; A 2.8.6; Select the elements which forms:
 - (1 mark) a double charged cation
 - (1 mark) a soluble carbonate
 - (1 mark) Which element has the shortest atomic radius?
- Study the scheme below and answer the questions that follow




(1 mark) a) Identify solid E

b)	Write an ionic equation for the reaction in step II that produces solid F	(1 mark)
Giv	re reason why phosphorus is stored under water	(1 mark)
C (289K and atmosphere pressure, graphite changes into diamond according to graphite) → C(diamond); ΔH = 2.9kJmol ⁻¹	7 8
In Log	the space provided, sketch a simple energy level diagram for the above change	e. (2 marks)
Ho	w would you obtain a sample of pure iodine from a mixture of iodine and lea	d sulphate? (2 marks)
10 jui a)	g of sodium hydrogen carbonate were dissolved in 20cm ³ of water in a boiling ce was then added dropwise with shaking until there was not further observate Explain the observation which was made in the boiling tube when the react progress What observation would have been made if the lemon juice had been added	ole change.
	turnings in a boiling tube? Give a reason	(1 mark
ar	mple solutions of salt were labelled as I, II, III and IV. The actual solutions, not lead nitrate, zinc sulphate, potassium chloride and calcium chloride. When aqueous sodium carbonate was added to each sample separately, a was formed in I, III and IV only. Identify solution II.	
-	was formed at 1, 111 data 1 v out, 1 and 1 v	
b)	When excess aqueous sodium hydroxide was added to each sample separat precipitate was formed in III only. Identify solution III.	ely, a white (1 mark
c)	When dilute sulphuric acid was added separately to each sample, a white p formed in solutions I and III only. Identify solution I.	recipitate was (1 mark
ur	weighed sample of crystalline sodium carbonate, (Na ₂ CO ₃ nH ₂ O) was heated it there was no further change in mass. The mass of the sample reduced by laculate the number of moles (n) of the water of crystallisation.	
	Ta = 23, O = 16, C = 12 H = 1)	(3 marks

CHVIIOIIIK	chiai chect of using	such matches in large numbers.	(2 marks
-	₹ <mark>%</mark>		
	at to		
	\$ CO.		
Describe a	a simple laboratory	experiment that can be used to distinguish betw	een sodium
mulahida a	A A Albamana	te.	(3 marks
e truit.	Kir		
*C28.70	33		
e win			
• \			• (0.13 bes. (0. • (7) • 4/7, (3.6) • (1)
		elow relates to elements in the same group of the	he period table.
Study it a	nd answer the quest		
	Element	Atomic size	
	G1	0.19	
	G2	0.23	
	G3	0.15	
Which ele	ment has the highes	st ionisation energy? Give a reason	(3 marks
		was heated with powdered carbon, the mixture	
carbon die		ed. When the experiment was repeated using th	
J, there w	oxide gas was forme as no apparent react	ed. When the experiment was repeated using th	e oxide of elemen
J, there w	oxide gas was forme as no apparent react	ed. When the experiment was repeated using th ion.	e oxide of elemen
J, there w	oxide gas was forme as no apparent react	ed. When the experiment was repeated using th ion.	e oxide of elemen
carbon die J, there wa a) Sugg	oxide gas was forme as no apparent react est one method that	ed. When the experiment was repeated using th ion.	e oxide of elemen
carbon die J, there wa a) Sugg	oxide gas was forme as no apparent react est one method that	ed. When the experiment was repeated using the ion. can be used to extract element J from its oxide.	e oxide of elemen
b) Arrar	oxide gas was formed as no apparent react test one method that the second secon	ed. When the experiment was repeated using the ion. can be used to extract element J from its oxide. J and carbon in the order of their decreasing reacted sulphuric acid was left in an open beaker in a	e oxide of elements (1 mark
b) Arrar When a sa	as no apparent react test one method that meet the elements H,	ed. When the experiment was repeated using the ion. can be used to extract element J from its oxide. J and carbon in the order of their decreasing reacted sulphuric acid was left in an open beaker in a polynomial or have increased slightly.	e oxide of elements (1 mark
b) Arrar When a sa	as no apparent react test one method that meet the elements H,	ed. When the experiment was repeated using the ion. can be used to extract element J from its oxide. J and carbon in the order of their decreasing reacted sulphuric acid was left in an open beaker in a	activity (1 mark
b) Arrar When a sa	as no apparent react test one method that meet the elements H,	ed. When the experiment was repeated using the ion. can be used to extract element J from its oxide. J and carbon in the order of their decreasing reacted sulphuric acid was left in an open beaker in a polynomial or have increased slightly.	activity (1 mark
b) Arrar When a sa days, the	as no apparent react test one method that test one method that make the elements H, ample of concentrate volume was found to property of concentrate property property of concentrate property property of concentrate property p	ed. When the experiment was repeated using the ion. can be used to extract element J from its oxide. J and carbon in the order of their decreasing reacted sulphuric acid was left in an open beaker in a polynomial or have increased slightly.	activity (1 mark

Connecting wire Connecting wire Magnesium metal Applain why it is necessary to connect the piece of magnesium metal to the iron bar State one cause of temporary hardness in water D) How does distillation remove hardness from water?	(3 marks) (1 mark) (2 marks)
a) State one cause of temporary hardness in water b) How does distillation remove hardness from water?	(1 mark)
State one cause of temporary hardness in water How does distillation remove hardness from water?	
	(2 marks)
- the CYLAY II a	
n the presence of U.V light, ethane gas undergoes substitution reaction with chlorine What is meant by the term: Substitution reaction?	(1 mark)
Cityo the structural family 141 City	
trobused of athere and the control of the organic product formed when eq	jual (2 marks)
Explain why burning magnesium continues to burn in a gas jar full of sulphur dioxide burning splint would be extinguished.	e while a (3 marks)
What observations would be made if hydrogen sulphide gas was bubbled through solution of zinc nitrate?	ha (1 mark)
Write an equation for the reaction that takes place in (a) above.	(1 mark)
1)	volumes of ethane and chlorine react together. cplain why burning magnesium continues to burn in a gas jar full of sulphur dioxiderning splint would be extinguished. What observations would be made if hydrogen sulphide gas was bubbled through solution of zinc nitrate?

21. The apparatus shown below was used to investigate the effect of carbon monoxide on copper



a) State the observation that was made in the combustion tube at the end of the experiment (1 mark)

- b) Write an equation for the reaction that took place in the combustion tube. (1 mark)
- c) Why is it necessary to burn the gas coming out of tube K? (1 mark)
- Explain why hydrogen forms compounds in which its oxidation state is either +1 or -1.
 (Atomic number of hydrogen is 1)

23. The table below shows the properties of substances K,L.M and N.

Substances	Reaction with-	Melting point	Conductivity	
	oxygen at 25°C	Plante Al VecTe PlantedEx	Solid	Molten
K	Unreactive	Low	Poor	Poor
L	Reactive			
M	Unreactive	High	Good	Good
N	Unreactive	Low	Good	Good

Select the substance which is likely to be:

a) Copper metal

(1 mark)

b) Magnesium chloride

(1 mark)

24. An element P has a relative atomic mass of 88. When a current of 0.5 amperes was passed through the fused chloride of P for 32 minutes and 10 seconds, 0.44g of P were deposited at the cathode. Determine the charge on an ion of P (1 Faraday = 96500 Coulombs) (3 marks) 25. The melting point of phosphorus trichloride is -91°C. while that of magnesium chloride is 715°C. In terms of structure and bonding, explain the difference in their melting points. (3 marks) 26. The pH of a sample of soil was found to be 5.0. An agricultural officer recommended the addition of calcium oxide in the soil 27. Hydrogen peroxide decomposes according to the equation given below. $H_2O_{2(j)} \to H_2O_{(j)} + \frac{1}{2}O_{2(g)}; \Delta H = -98kJ$ 8.5 of hydrogen peroxide contained in 100cm3 of solution with water were completely

decomposed.

Determine the rise in temperature due to the reaction.

(Specific heat capacity of water = $4.2 \text{Jg}^{-1} \text{K}^{-1}$, density of water = 1g/cm^3 O = 16 H = 1)

(3 marks)