



THE UNIVERSITY
of LIVERPOOL

SUMMER 2000 EXAMINATIONS

Degree of Bachelor of Engineering : Year 2
Degree of Bachelor of Engineering : Year 3
Degree of Master of Engineering : Year 2

ENGINEERING MATHEMATICS II

TIME ALLOWED : Two Hours

INSTRUCTIONS TO CANDIDATES

Full marks can be obtained for complete answers to FOUR questions.
Only the best FOUR answers will be counted.



THE UNIVERSITY
of LIVERPOOL

1. (a) Evaluate the determinant

$$\begin{vmatrix} 2 & 3 & 0 & 1 \\ 1 & 2 & 3 & -1 \\ 1 & 2 & 0 & 1 \\ 2 & -2 & 1 & 2 \end{vmatrix}$$

(10 marks)

- (b) Use Cramer's rule to solve the system

$$\begin{aligned} x + y + 3z &= 6 \\ 2x + 2y + \beta z &= -2 + 4\beta \\ x - y + 2z &= 2 \end{aligned}$$

when $\beta = 1$.

(11 marks)

- (c) At what β does this system have no solutions?

(4 marks)



THE UNIVERSITY
of LIVERPOOL

2. (a) **A** and **B** are two square matrices and $\det(\mathbf{A}) \neq 0$. Show that a matrix **C** satisfying

$$\mathbf{CA} = \mathbf{AB}$$

is given by

$$\mathbf{C} = \mathbf{ABA}^{-1}.$$

(2 marks)

- (b) Further, show that a matrix **D** satisfying

$$\mathbf{DA} = \mathbf{AB}^2$$

is given by

$$\mathbf{D} = \mathbf{C}^2.$$

(5 marks)

- (c) Find $\mathbf{C} = \mathbf{ABA}^{-1}$, if

$$\mathbf{A} = \begin{bmatrix} 1 & 3 & 1 \\ 0 & 2 & 1 \\ -1 & -2 & 0 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

(14 marks)

- (d) Verify the above result by showing that $\mathbf{CA} = \mathbf{AB}$

(4 marks)



THE UNIVERSITY
of LIVERPOOL

3. (a) Show that $\lambda = 1$ is one eigenvalue of the matrix

$$\mathbf{A} = \begin{bmatrix} -1 & -1 & 1 \\ 2 & -1 & 2 \\ 2 & 1 & 0 \end{bmatrix}$$

and find the remaining eigenvalues.

(8 marks)

- (b) Find corresponding eigenvectors.

(8 marks)

- (c) Write the system of differential equations for $x(t)$, $y(t)$ and $z(t)$, corresponding to the matrix differential equation

$$\frac{d\mathbf{u}}{dt} = \mathbf{A}\mathbf{u} \quad \text{where} \quad \mathbf{u}(t) = \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

(2 marks)

- (d) Using the above results on the eigenvalues and eigenvectors of \mathbf{A} , find the general solution of this system. Check your solution.

(7 marks)



THE UNIVERSITY
of LIVERPOOL

4. (a) State the linear approximation to a function $f(x, y)$ in the neighbourhood of the point (x_0, y_0) .

(3 marks)

- (b) The function $f(x, y)$ is defined as

$$f(x, y) = e^{x^2 - y^2}.$$

Find the partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$. Hence find the linear approximation to this function in the neighbourhood of the points

- $x_0 = 1, y_0 = 0$;
- $x_0 = 0, y_0 = 1$.

(10 marks)

- (c) Find the second derivatives

$$\frac{\partial^2 f}{\partial x^2} \text{ and } \frac{\partial^2 f}{\partial y^2}$$

for the function $f(x, y)$ defined above, and hence evaluate

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2},$$

stating your result in its simplest form.

(12 marks)



THE UNIVERSITY
of LIVERPOOL

5. (a) Briefly show how to derive the explicit finite difference scheme:

$$y_{n+1} = y_n + \frac{1}{2}\Delta_n + \frac{1}{2}hf(x_n + h, y_n + \Delta_n)$$

where

$$\Delta_n = hf(x_n, y_n)$$

for the numerical solution of the nonlinear differential equation

$$dy/dx = f(x, y)$$

with initial conditions

$$y(x_0) = y_0$$

(It is **not** necessary to discuss the error in this approximation).

(12 marks)

- (b) Use this scheme, with a step length $h = 0.025$, to calculate approximately $y(2.1)$ when

$$f(x, y) = y^2 - \frac{1}{x^2} \quad \text{and} \quad y(2) = 1.$$

(13 marks)